

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

**A Thesis Submitted for the Degree of PhD at the University of Warwick**

<http://go.warwick.ac.uk/wrap/74317>

This thesis is made available online and is protected by original copyright.

Please scroll down to view the document itself.

Please refer to the repository record for this item for information to help you to cite it. Our policy information is available from the repository home page.

# **Equivariant cobordism and K-theory**

**Mohamed Abdel Moniem Ismail**

**Warwick University**

**1971**

**PhD**

# CONTENTS

## INTRODUCTION

### Chapter 1 : Equivariant Cohomology Theories and Cobordism Theory:

	Page
1. G-spectra	1
2. Continuity	5
3. $U_G^*$ -theory	8
4(A) Multiplication	12
4(B) The Thom Isomorphism	16
4(C) The isomorphism $F(G, H)$	20
5(a) Spectral Sequences	23
5(b) Free Group Action	26
6. Multiplicative Equivariant Theories	28

### Chapter 2 : The Equivariant Cobordism of Projective Spaces

1. The Conner-Floyd Map $\mu$	1
2. Some Remarks	6
3. Localization	8
4. $U_G^*(-)[\wedge^{-1}]$ of Projective Spaces	11

### Chapter 3 : Applications

1. Characteristic Classes	16
2. An Embedding of $K_G(X, A)$	20
3. $U_G^*(-)[\wedge^{-1}]$ of Grassmannians	23
4. Equivariant K-theory and Cobordism (G abelian)	28

### Chapter 4 : The Relation of $K_G$ - theory to Cobordism

1. The Relation of $U_G^*(-)$ with Bordism	30
2. A Gysin homomorphism	34
3. $Td([G/T])$	37
4. The Main Theorem	39

# INTRODUCTION :

This thesis is mainly a study of the equivariant unitary cobordism  $U_G^*$ ,  $G$  a compact Lie group. It is obtained in Chapter 1 as an example of an equivariant cohomology theory associated to a  $G$ -spectrum. In addition to discussing a number of results needed in the rest of the thesis, we set up the  $U_G^*$ -spectral sequence and use it to prove that given a compact  $G \times G'$ -space  $X$  such that  $G$  acts freely on  $X$ , there is a natural multiplicative isomorphism  $pr^* : U_{G'}^*(X/G) \longrightarrow U_{G \times G'}^*(X)$ .

In Chapter 2, we define a localized cobordism theory  $U_G^*[\wedge^{-1}]$ . Given an  $n$ -dimensional  $G$ -vector bundle  $E$  over a compact  $G$ -space  $X$ , ( $G$  abelian), it is proved that  $U_G^*(P(E))[\wedge^{-1}]$  is freely generated over  $U_G^*(X)[\wedge^{-1}]$  by  $1, \rho, \dots, \rho^{n-1}$  for some natural element  $\rho \in U_G^2(P(E))[\wedge^{-1}]$ . This is exploited in Chapter 3 to define  $U_G^*[\wedge^{-1}]$ -characteristic classes, to give a partial result about  $U_G^*[\wedge^{-1}]$ -theory of Grassmannians, and prove that (for  $G$  abelian)

$$U_G^*(X, A) \underset{U_G}{\otimes}^* RG \overset{\vee}{=} K_G^*(X, A) \dots \dots \dots (*)$$

for  $X$  a locally compact  $G$ -space and  $A$  a closed  $G$ -subspace of  $X$ .

Chapter 4 deals with the proof of this latter result for non-abelian  $G$ . For connected  $G$  with maximal torus  $T$  a Gysin homomorphism  $\tau! : U_T^*(X) \longrightarrow U_G^*(X)$  ( $X$  a compact  $G$ -space) is defined. Composed with the restriction homomorphism  $U_G^*(-) \longrightarrow U_T^*(-)$ , the result is multiplication by the bordism class  $[G/T]$ .



Well-known results of Atiyah, Segal and Singer are then used to verify that the equivariant Todd genus of  $G/T$  is equal to  $1 \otimes \text{ERG}$ . This leads quite easily to the main theorem (the isomorphism (\*) for general  $G$ ).

I am most grateful to my supervisor Dr. Luke Hodgkin for his help, his encouragement, and his suggestions throughout the last three years. I would like to thank too Dr. T. tom Dieck for his useful suggestions.

§ 1.1 : G-spectra:-

Let  $G$  be a compact Lie group. By a  $G$ -module, we shall mean finite-dimensional complex representation of  $G$ . Let  $(\underline{c}_1)$  be the category whose objects are pairs of spaces  $(X, A)$ ,  $X$  a locally compact  $G$ -space and  $A$  a closed  $G$ -subspace of  $X$  and whose morphisms are proper  $G$ -maps of pairs. We shall regard a locally compact  $G$ -space  $X$  as an object of  $(\underline{c}_1)$  by identifying it with  $(X, \emptyset)$  ( $\emptyset$  is the empty set). Let  $\text{ab}$  be the category of abelian groups, and homomorphisms. An equivariant cohomology theory  $h_G^* = (h_G^n : n \in \mathbb{Z})$  on  $(\underline{c}_1)$  is defined to be a sequence of contravariant functors:  $h_G^n : (\underline{c}_1) \longrightarrow \text{Ab}$  and a sequence of natural transformations  $\partial^n : h_G^n \longrightarrow (A) \longrightarrow h_G^{n+1}(X, A) \forall (X, A) \in (\underline{c}_1)$  such that:

(a1) if  $f \simeq f^1 : (X, A) \longrightarrow (X^1, A^1)$  ( $\simeq$  : means  $G$ -homotopic by proper  $G$ -maps), then  $h_G^n(f) = h_G^n(f^1)$

(a2) if  $i: X, A \hookrightarrow X, A$  ,  $j: (A, \emptyset) \longrightarrow (X, \emptyset)$  are the inclusions, then the sequence

$$\dots h_G^{n-1}(A) \xrightarrow{\partial^{n-1}} h_G^n(X, A) \xrightarrow{h_G^n(j)} h_G^n(X) \xrightarrow{h_G^n(i)} h_G^n(A) \xrightarrow{\partial^n} \dots$$

is exact

(a3) every relative  $G$ -homeomorphism  $(X, A) \xrightarrow{f} (X^1, A^1)$  of objects of  $(\underline{c}_1)$ , (i.e.  $X/A \longrightarrow X^1/A^1$  is a  $G$ -homeomorphism) induces an isomorphism  $h_G^*(f)$ ,

(a4)  $h_G^*$  is additive i.e. for any family of spaces  $(X_\alpha)$  in  $(\mathcal{C}_1)$ , the natural homomorphism:

$$h_G^* (\coprod_\alpha X_\alpha) \longrightarrow \prod_\alpha h_G^* (X_\alpha)$$

is an isomorphism.

Notation: If  $f$  is a morphism in  $(\mathcal{C}_1)$ , we shall write  $f^*$  for  $h_G^n(f)$  as usual.

As in the ordinary case ([12]), there is a natural 1-1 correspondence between theories  $(h_G^*)$  and reduced equivariant cohomology theories  $(\tilde{h}_G^*)$  defined on the category of compact  $G$ -spaces with base point.

A  $G$ -spectrum consists of:

(i) a sequence  $(Y_k : k = 0, \pm 1, \pm 2, \dots)$  of  $G$ -spaces with base points

(ii)  $\forall$   $G$ -module  $V$ , a base point preserving  $G$ -map:  $V^+ \wedge Y_k \rightarrow Y_{|V|+k}$

where  $|V|$  = dimension of  $V$  over  $\mathbb{C}$  and  $X^+$  denotes the one point compactification of the locally compact  $G$ -space  $X$ . The diagram:

$$\begin{array}{ccc} (V \oplus W)^+ \wedge Y_k & = & V^+ \wedge (W^+ \wedge Y_k) \longrightarrow W^+ \wedge Y_{|V|+k} \\ & \searrow & \downarrow \\ & & Y_{|V|+|W|+k} \end{array}$$

is  $G$ -homotopy commutative for all  $V \oplus W$ .

The importance of  $G$ -spectra lies in the fact that each defines an equivariant cohomology theory as was shown by G. Whitehead

[24] for  $G=e$ . For let  $Y_G = \{Y_k\}$  be a  $G$ -spectrum and let  $X$  be a compact  $G$ -space with base point. Let  $\Delta_G$  be the set of all  $G$ -modules, with the partial order relation:  $V \leq W$  if  $\exists f$  a

$G$ -module  $V^1$  s.t.  $V \oplus V^1 = W$ . This makes  $\Delta_G$  into a directed set. Given  $V \in \Delta_G$  &  $n \in \mathbb{Z}$ , define

$$B_n^V = [V^+ \wedge X, Y_{|V|+n}]_G^0$$

= the set of  $G$ -homotopy classes of based  $G$ -maps from

$$V^+ \wedge X \rightarrow Y_{|V|+n}. \text{ Given } V \oplus V^1 = W, \text{ define}$$

$$f_V^W : B_n^V \rightarrow B_n^W$$

by :  $f_V^W (V^+ \wedge X \xrightarrow{t} Y_{|V|+n}) = \text{the composition}$

$$(V \oplus V^1)^+ \wedge X = V^{1+} \wedge (V^+ \wedge X) \xrightarrow{1 \wedge t} V^{1+} \wedge Y_{|V|+n} \rightarrow Y_{|W|+n}$$

where the latter map comes from the definition of

$\mathcal{Q}_G = \{Y_k\}$ . The collection  $\{B_n^V, f_V^W\}_{\Delta_G}$  is a direct system and if  $W = V \oplus C$ ,  $W^+ = V^+ \wedge S^2$ , so  $[W^+ \wedge X, \dots]_G^0$  is an abelian gp.

Define (1.1.1):

$$\tilde{\mathcal{Q}}_G^{2n}(X) = \varinjlim [V^+ \wedge X, Y_{|V|+n}]_G^0 \quad - \text{it is}$$

naturally an abelian group. Define :

$$\tilde{\mathcal{Q}}_G^{2n-1}(X) = \tilde{\mathcal{Q}}_G^{2n}(SX).$$

Given a closed  $G$ -subspace  $A$  of a compact  $G$ -space  $X$  with base point  $x_0 \in A$ , define :

$$\tilde{\mathcal{Q}}_G^n(X, A) = \tilde{\mathcal{Q}}_G^n(X \cup_A CA)$$

where  $CA$  is the reduced cone on  $A$ . As in [11] we can define a

map  $\sigma_n : \tilde{\mathcal{Q}}_G^n(X) \rightarrow \tilde{\mathcal{Q}}_G^{n+1}(SX)$  as follows: suppose  $n = 2k$

and let  $x \in \tilde{\mathcal{Q}}_G^{2k}(X)$  be represented by  $f: V^+ \wedge X \rightarrow Y_{|V|+k}$ .

Assign to  $x$  the element  $\sigma(x)$  represented by the composition:

$$V^+ \wedge S(SX) = S^2 \wedge (V^+ \wedge X) \xrightarrow{1 \wedge f} S^2 \wedge Y_{1V1+k} \longrightarrow Y_{1V1+k+1}$$

where the last map comes from  $Y_G$ . From the definitions, it is immediate that  $\sigma$  is an isomorphism.

Proposition 1.1.2:

$\tilde{Y}_G^* = \{ \tilde{Y}_G^h, \sigma \}$  is a reduced equivariant cohomology theory on the category of compact  $G$ -spaces with base point. Hence there is a corresponding equivariant cohomology theory  $Y_G^*$  defined on the category  $(\underline{C}_1)$ .

Proof:-

The first part follows by using the method of Dyer [12] (p.9-13).

The analogue of lemma 3, namely that

$$V^+ \wedge (CA \cup X) = C(V^+ \wedge A) \cup (V^+ \wedge X)$$

for a closed  $G$ -subspace  $A$  of a compact  $G$ -space  $X$  with base point  $x_0 \in A$  follows by noticing that each of the spaces in question is obtained from  $V^+ \times (I \times A) \cup V^+ \times X$  by making the same identifications. The rest of the proof applies equally well here.

The 2nd part of the Proposition 1.1.2 follows from the first part (Dyer [12])

Q.E.D.

Let us recall that as in the ordinary case,  $G=e$ , we define  $Y_G^*$  as follows:

Definition 1.1.3:

Let  $X$  be a locally compact  $G$ -space and  $A$  a closed  $G$ -subspace

of  $X$ . Define  $y_G^n(X) = \tilde{y}_G^n(X^+)$ ,

$$y_G^n(X, A) = \tilde{y}_G^n(X^+, A^+).$$

§ 1.2: Continuity:

Define 1.2.1 :

A family  $F$  of pairs of closed  $G$ -subspaces of a locally compact  $G$ -space  $X$  is said to be a filtering family if  $(L, B), (L', B') \in F$ ,  $\Rightarrow \exists (L'', B'') \in F$  s.t.  $(L'', B'') \subset (L \cap L', B \cap B')$

Proposition 1.2.2 :

Let  $y_G^*$  be the equivariant cohomology theory associated to a  $G$ -spectrum  $y_G = \{Y_k\}_{k \in \mathbb{Z}}$ . Suppose each  $Y_k$  is the union of compact  $G$ -subspaces each of which can be embedded in a  $G$ -module as an equivariant neighbourhood retract. Then given a filtering family  $F$  of closed  $G$ -subspaces of a locally compact  $G$ -space  $X$ , the natural homomorphism.

$$\theta : \lim_{\substack{\longrightarrow \\ F}} y_G^*(L, B) \rightarrow y_G^* \left( \bigcap_{(L, B) \in F} L, \bigcap_{(L, B) \in F} B \right)$$

is an isomorphism. In particular, if  $F$  is the family of closed  $G$ -neighbourhoods of a closed  $G$ -subspace  $A$  of  $X$ , then :

$$\theta : \lim_{\longrightarrow} y_G^*(F) \xrightarrow{\cong} y_G^*(A).$$

Proof:-

Since  $\bigcap (L^+ \cup B^+ \cup CB^+) = \bigcap L^+ \cup \bigcap B^+ \cup \bigcap CB^+ \subset \bigcap B^+$ , we need only show that  $\varinjlim \tilde{y}_G^* (L) \xrightarrow{\cong} \tilde{y}_G^* (\bigcap L)$ ,

when all the  $L$ 's are compact and have a common base point.

Put  $A = \bigcap L$  and let  $x \in \tilde{y}_G^* (A)$  be represented by the map  $f : V^+ \wedge A \rightarrow Y_k$ . Hence  $f(V^+ \wedge A) \in M$ , a  $G$ -subspace of  $Y_k$

s.t.  $M$  can be embedded in a  $G$ -module  $W$  by an embedding

$j : M \rightarrow W$ . In addition  $\exists$  a neighbourhood  $N$  of  $M$  in  $W$  and

a retraction  $r : N \rightarrow M$ . By the Tietze extension theorem,

extend the map  $j \circ f : V^+ \wedge A \rightarrow W$  to the whole of  $V^+ \wedge X$  thus

getting a map  $h : V^+ \wedge X \rightarrow W$ . Hence we get a map:

$\phi : (V^+ \wedge X) \rightarrow W$  defined by  $(g, y) \rightarrow g(h(g^{-1}(y)))$ . Integrating

over  $G$  (Pontjagin [19]) we obtain an equivariant map

$h_1 : V^+ \wedge X \rightarrow W$  s.t.  $j \circ f = h_1|_{V^+ \wedge A} \rightarrow W$ . By continuity

of  $h_1$ ,  $\exists$  a  $G$ -neighbourhood  $T$  of  $A$  s.t.  $h_1(V^+ \wedge T) \subset N$ . We now

have a  $G$ -map  $V^+ \wedge T \rightarrow Y_k$  defined by the composition:

$$V^+ \wedge T \xrightarrow{h_1} N \xrightarrow{r} M \xrightarrow{j} Y_k,$$

which restricted to  $V^+ \wedge A$  is  $= f$ . Since  $T$  contains a closed

$G$ -neighbourhood of  $A$  (by compactness), it follows that

$$e : \varinjlim \tilde{y}_G^* (L) \rightarrow \tilde{y}_G^* (A)$$

is an epimorphism.

Suppose now  $x, y \in \tilde{y}_G^* (L)$ , represented by:

$$f : W^+ \wedge L \rightarrow Y_k, \quad g : W^+ \wedge L \rightarrow Y_k$$

satisfy :  $i^*(x) = i^*(y)$

where  $i : A \hookrightarrow L$  is inclusion. This means  $\exists$  a  $G$ -module  $V$  s.t.

the composition :  $V^+ \wedge (W^+ \wedge A) \xrightarrow{1 \wedge f} V^+ \wedge Y_k \xrightarrow{(1)} Y_{|V|+k}$

is  $G$ -homotopic to the composition:

$$V^+ \wedge (W^+ \wedge A) \xrightarrow{1 \wedge g} V^+ \wedge Y_k \xrightarrow{(1)} Y_{|V|+k}$$

(here (1) is the given map of the spectrum) i.e. there is

$$\text{a } G\text{-map } (V^+ \wedge W^+ \wedge A) \times I \cup (V^+ \wedge W^+ \wedge L) \times dI \xrightarrow{f^*} Y_{|V|+k} \\ (V^+ \wedge W^+ \wedge A) \times \partial I$$

Now  $(V^+ \wedge W^+ \wedge A) \times I \cup (V^+ \wedge W^+ \wedge L) \times dI$  is a closed  $G$ -subspace of  $(V^+ \wedge W^+ \wedge L) \times I$ . By the same argument used above,

$\exists$  a closed  $G$ -neighbourhood  $N^*$  of

$$(V^+ \wedge W^+ \wedge A) \times I \cup (V^+ \wedge W^+ \wedge L) \times dI \text{ and a } G\text{-map} \\ (V^+ \wedge W^+ \wedge A) \times dI$$

$$\tilde{f}^* : N^* \rightarrow Y_{|V|+k}$$

extending  $f^* : (V^+ \wedge W^+ \wedge A) \times I \cup (V^+ \wedge W^+ \wedge L) \times dI \rightarrow Y_{|V|+k} \\ (V^+ \wedge W^+ \wedge A) \times dI$

It is very easy to see that  $N^* \supset (V^+ \wedge W^+ \wedge \dot{L}) \times I$  for some  $L' \in L$ .

Therefore  $j^* \circ \theta = j^*$  where  $j : L' \hookrightarrow L$  is the inclusion

i.e.  $\theta$  is a monomorphism.

Q.E.D.

#### Remark:

The method above can also be used to prove continuity of  $K_G$ -theory (Cf Segal [20]). The only non-trivial point is extending a



$G$ -vector bundle  $E$  on  $A$  to a closed  $G$ -neighbourhood of  $A$  in  $X$  (all spaces being compact). If  $A$  is connected, then  $E$  is classified by a  $G$ -map

$$f : A \rightarrow G_k(V)$$

where  $k = \dim E$ , and  $G_k(V)$  is the Grassmann Manifold of  $(k - \dim \mathbb{C})$  subspaces of a  $G$ -module  $V$ .  $G_k(V)$  can be embedded in a  $G$ -module  $W$  (Palais [18]) since it is a compact differentiable manifold and it is then a  $G$ -neighbourhood retract (the normal bundle to  $G_k(V) = \text{tubular neighbourhood of } G_k(V)$ ). Hence one can extend  $E$  to a neighbourhood of  $A$ . One deals with the general case by using the fact that  $S^2 \wedge A$  is connected.

### § 1.3 : The equivariant unitary cobordism theory $U_G^*(-)$ :-

#### 1.3.1: The Thorm Spectrum $MU_*(G)$ :-

Let  $\{V_j : j \in J\}$  be a complete set of irreducible inequivalent representations of the compact Lie group  $G$  ( $J$  is finite or countable according as  $G$  is finite or not). Define: the universal  $G$ -module  $V^\infty$  by:

$$V^\infty = \lim_{\substack{m \rightarrow \infty \\ m \in \mathbb{Z}}} \sum_{j \in J} V_j$$

$$\text{Let } \gamma_k^G = (E_k(G) \xrightarrow{\pi} B_k(G))$$

be the universal  $k$ -dimensional  $G$ -vector bundle i.e.

$E_k(G) = \{(K, x) : K \text{ a } k\text{-dim } \mathbb{C} \text{ plane through the origin in } V^\infty, \text{ and } x \in K\}$ ,  $B_k(G) = \{K : K \text{ a } k\text{-plane through the origin in } V^\infty\}$  &  $\pi : E_k(G) \rightarrow B_k(G)$  is projection onto the first factor.

The reason these bundles are called universal is :

Proposition 1.3.1 :

Let  $X$  be a compact  $G$ -space. The assignment:

$$[f] \rightarrow [f^* (\nu_k^G)]$$

defines a natural 1-1 correspondence between the set of  $G$ -homotopy classes of  $G$ -maps from  $X$  to  $B_k(G)$  and the set of isomorphism classes of  $k$ -dim.  $G$ -vector bundles on  $X$ .

Proof:

Denote as usual the set of  $G$ -homotopy classes of  $G$ -maps from  $X$  to  $B_k(G)$  by  $[X, B_k(G)]_G$  and the set of isomorphism classes of  $k$ -dim.  $G$ -vector bundles on  $X$  by  $\text{Vect}_k^G(X)$ .

By Proposition 1.3 (Segal [20]), if  $f' \simeq f : X \rightarrow B_k(G)$ , then  $f'^* (\nu_k^G) \cong f^* (\nu_k^G)$ . Hence we have a well-defined map:

$$[X, B_k(G)]_G \longrightarrow \text{Vect}_k^G(X)$$

On the other hand, if  $E$  is a  $k$ -dim.  $G$ -vector bundle on  $X$ , there is a  $G$ -vector bundle  $E^\perp$  on  $X$  s.t.

$E \oplus E^\perp$  is trivial,  $E \oplus E^\perp = \underline{V} = X \times V$  for some  $G$ -module  $V$  (Proposition 2.4 Segal [20]). Hence there is an epimorphism  $\psi : X \times V \rightarrow E$  inducing a  $G$ -map :

$f : X \rightarrow G_k(V)$  (the Grassmann manifold of  $k$ -dim. subspaces of  $V$ ) by assigning to  $x \in X$ ,  $f(x) = V/Ker \psi_x$ . Proceeding as in Atiyah ([1] P. 29), we can prove that the homotopy class of the  $G$ -map :

$$X \xrightarrow{\tilde{f}} B_k(G)$$

given by the composition  $X \xrightarrow{f} G_k(V) \subset B_k(G)$ , does not depend

on  $V$  or the epimorphism  $\Psi$  we choose. So we construct an inverse map  $\text{Vect}_k^G(X) \rightarrow [X, B_k(G)]_G$  by :

$$[E] \rightarrow [\bar{f}] \quad \text{Q.E.D.}$$

By analogy with the ordinary case,  $G=e$ , where the  $B_k(G)$  are the infinite - dimensional Grassmannians, we can see that  $B_k(G)$  is paracompact and thus  $\gamma_k^G$  admits an invariant metric (Milnor [15], Husemoller [14]). Define the universal k-dimensional Thom space  $M_k(G)$  by

$$M_k(G) = D(E_k(G)) / S(E_k(G)) = \frac{\text{the unit ball bundle of } E_k G}{\text{the unit sphere bundle of } E_k G}$$

w.r.t. the above metric. So  $M_k(G)$  has a natural base point.

Moreover,  $\forall$   $G$ -module  $V$ , we have a natural  $G$ -homotopy class of maps:

$$V^+ \wedge M_k(G) \longrightarrow M_{|V|+k}(G)$$

namely the one induced by the classifying map of the bundle :

$$V \times E_k(G) \longrightarrow \text{pt.} \times B_k(G)$$

(Proposition 1.3.1).  $MU_*(G) = \{M_k(G)\}$  defines a  $G$ -spectrum, called the equivariant. Thom spectrum.

Hence, by § 1.1, the following definitions make sense.

Definition 1.3.2 :

- (i) Given a compact  $G$ -space  $X$  with base point, and a closed  $G$ -subspace  $A$  of  $X$ , containing the base point,

$$\text{define } \tilde{U}_G^{2n}(X) = \varinjlim [V^+ \wedge X, M_{|V|+n}(G)]_G^0,$$

$$\tilde{U}_G^{2n-1}(X) = \tilde{U}_G^{2n}(SX) \quad \&$$

$$\tilde{U}_G^n(X, A) = \varinjlim_A \tilde{U}_G^n(X \cup CA).$$

(ii) Given a locally compact G-space X, and a closed G-subspace A of X, define

$$U_G^n(X) = \check{U}_G^n(X^+) \text{ \& } U_G^n(X, A) = \check{U}_G^n(X^+, A^+)$$

It follows from Proposition 1.1.2

Cor 1.3.3:

$U_G^* = \{U_G^n\}$  is an equivariant cohomology theory on the category  $(\underline{C}_1)$ .

COR 1.3.4:

$U_G^*$  is continuous i.e.  $\forall$  filtering family F of closed G-subspaces of a locally compact G-space X, the natural homomorphism:

$$\varinjlim_F U_G^*(L, B) \longrightarrow U_G^* \left( \bigcap_{(L, B) \in F} L, \bigcap_{(L, B) \in F} B \right)$$

is an isomorphism.

Proof:

By Proposition 1.2.2, it is enough to show that for all k,  $M_k(G)$  is the union of compact G-subspaces each of which can be embedded in a G-module as a G-neighbourhood retract. Consider a G-module V. Let  $M_k(V)$  be the Thom-space of the standard k-dim  $\mathbb{C}$ -G-vector bundle  $E_k(V)$  on  $G_k(V) =$  the Grassmann manifold of k-dim  $\mathbb{C}$ -subspaces of V.  $M_k(V)$  can be naturally identified with a subspace of  $M_k(G)$  and as V varies over the set of G-modules,  $\Delta G$ , the union of these subspaces is equal to  $M_k(G)$ . Since  $G_k(V)$  is a compact differentiable G-manifold, the disc bundle  $D(E_k(V))$  is a compact differentiable G-manifold. According to Palais [18] we can embed  $D(E_k(V))$  in

a  $G$ -module  $W$  by an embedding  $f$  say. This in turn induces an embedding  $f^*$  of  $M_k(V)$  in  $W \oplus \mathbb{C}$  defined by :

$$f^*(x) = ((1 - \|x\|) f(x), \|x\|) \quad (\dagger)$$

Let  $S(E_k(V))$  be the sphere bundle of  $E_k(V)$ . The map  $f^* : \text{int } D = D(E_k(V)) \setminus S(E_k(V)) \longrightarrow W \oplus \mathbb{C}$  assigning to  $x$  the element  $((1 - \|x\|) f(x), \|x\|)$  is an embedding of the  $G$ -manifold  $\text{int } D$  in  $W \oplus \mathbb{C}$ . We can identify the normal bundle to this manifold in  $W \oplus \mathbb{C}$  with a neighbourhood  $N$  of it in  $W \oplus \mathbb{C}$ . Let  $N^*$  be a small ball with centre the point  $(0,1)$  in  $W \oplus \mathbb{C}$ .  $NUN^1$  is a neighbourhood of  $f^*(M_k(V))$ , and  $N$  retracts onto  $f^*(\text{int } D)$  while  $N^1$  retracts onto  $(0,1)$ . We can now find a closed  $G$ -neighbourhood of  $f^*(M_k(V))$  which is a subspace of  $NUN^1$ , and retracts onto  $f^*(M_k(V))$ . This completes the proof.

Q.E.D.

#### §1.4 (A): Multiplication and some functorial Properties:-

For every pair of integers  $(m,n) \in \mathbb{Z}^+ \times \mathbb{Z}^+$ , there is a unique  $G$ -homotopy class of maps:

$$M_m(G) \wedge M_n(G) \xrightarrow{(m,n)} M_{m+n}(G)$$

It is the one induced by a classifying map of the bundle

$\gamma_m^G \times \gamma_n^G$ . This in turn induces a multiplication on  $\tilde{U}_G^*(X) \forall$

compact  $G$ -space  $X$  with base point defined as follows: Suppose

$x, y \in \tilde{U}_G^*(X)$  are represented by the  $G$ -maps  $V^+ \wedge X \xrightarrow{u} M_k(G)$

and  $W^+ \wedge X \xrightarrow{v} M_l(G)$ . Define  $xy \in \tilde{U}_G^*(X)$  to be the element

represented by the composition:

$$(\dagger) \quad \text{Read : } ((1 - \|x\|) f(x), \|x\|) .$$

$$(V \otimes W)^+ \wedge X \xrightarrow{1 \wedge \Delta} (V \otimes W)^+ \wedge X \wedge X = (V^+ \wedge X) \wedge (W^+ \wedge X) \xrightarrow{u \wedge v} M_k(G) \wedge M_l(G) \\ \downarrow (k,l) \\ M_{k+l}(G)$$

where  $\Delta : X \rightarrow X \times X$  is the diagonal map. This gives  $U_G^*(X) = \tilde{U}_G^*(X^+)$  a ring structure  $\forall$  locally compact  $G$ -space  $X$ . In particular, it makes  $\tilde{U}_G^* = U_G^*(\text{point})$  a ring with unit  $1$  represented by:

$$\mathbb{C}^+ \longrightarrow M_1(G) \quad (\text{trivial action on } \mathbb{C})$$

induced by the classifying map of the bundle  $\mathbb{C} \rightarrow \text{pt.}$

Let  $\alpha : G' \rightarrow G$  be a homomorphism of Lie groups. Let  $X$  be a compact  $G$ -space with base point,  $Y$  a compact  $G'$ -space with base point and  $S : Y \rightarrow X$  a map such that

$$s(g'y) = \alpha(g') s(y) \quad \forall g' \in G', y \in Y \quad \dots (*)$$

We shall call such a map  $S$  an  $\alpha$ -map for any given

$\alpha : G' \rightarrow G$ . Let  $x \in U_G^*(X)$  be represented by :

$V^+ \wedge X \xrightarrow{f} M_k(G)$ . Regard  $V$  as a  $G'$ -module via:

$g'(v) = \alpha(g')v$ . In the same way, we regard  $\gamma_k^G$  as a

$G'$ -vector bundle. So we get a unique  $G'$ -homotopy class:

$$M_k(G) \xrightarrow{\nabla} M_k(G')$$

Hence we can define a multiplicative homomorphism:

$$R\alpha : \tilde{U}_G^*(X) \longrightarrow \tilde{U}_{G'}^*(Y)$$

by:  $R\alpha(x)$  is the element of  $\tilde{U}_{G'}^*(Y)$  represented by the composition:

$$V^+ \wedge Y \xrightarrow{1 \wedge S} V^+ \wedge X \xrightarrow{f} M_k(G) \xrightarrow{\nabla} M_k(G').$$

Proposition 1.4.1 :

Let  $\alpha: G' \rightarrow G$ ,  $\alpha^1: G'' \rightarrow G'$  be homomorphisms of Lie groups. Suppose  $X$  is a compact based  $G$ -space,  $Y$  a compact based  $G'$ -space and  $Z$  a compact based  $G''$ -space. Suppose  $S: Y \rightarrow X$  is an  $\alpha$ -map,  $s': Z \rightarrow Y$  an  $\alpha'$ -map. The diagram:

$$\begin{array}{ccc} \tilde{U}_G^*(X) & \xrightarrow{R\alpha} & \tilde{U}_{G'}^*(Y) \\ & \searrow R(\alpha\alpha') & \swarrow R\alpha' \\ & \tilde{U}_{G''}^*(Z) & \end{array}$$

is commutative.

The proof is straightforward.

If  $\alpha: H \rightarrow G$  is the inclusion of a subgroup of  $G$  &  $S = \text{id}: X \rightarrow X$ , we shall call  $R$  the restriction homomorphism and denote it by  $r$ .

Let  $U^* \xrightarrow{r_1} U_G^*$  be the <sup>natural</sup> ~~restriction~~ homomorphism induced by the inclusion  $\{e\} \xrightarrow{\uparrow} G$ . According to Thom [23], we can identify  $U^*$  with the bordism ring of unitary manifolds  $\mathcal{U}_*$  by a natural identification:

$$i: \mathcal{U}_* \rightarrow U^*.$$

Suppose now  $X$  is a compact  $G$ -space. Let  $U_G^* \xrightarrow{r_2} U_G^*(X)$  be the natural map induced by the map  $X \rightarrow \text{point}$ . We can regard  $U_G^*(X)$  as a  $\mathcal{U}_*$ -module by means of the homomorphism:

$$\mathcal{U}_* \xrightarrow{i} U^* \xrightarrow{r_1} U_G^* \xrightarrow{r_2} U_G^*(X)$$

Proposition 1.4.2:

$U_G^*(X)$  is an algebra over  $\mathcal{U}^*$  for all compact  $G$ -spaces  $X$ .

† Give everything trivial  $G$ -action.

Lemma 1.4.3.:

Suppose  $X$  is a locally compact trivial  $G$ -space. The restriction

$$\text{homomorphism : } U_G^*(X) \xrightarrow{r} U^*(X)$$

has a natural right inverse :

$$r_1 : U^*(X) \longrightarrow U_G^*(X)$$

Proof:-

Let  $x \in U^*(X)$  be represented by :

$$S^{2k} \wedge X^+ \xrightarrow{f} M_{k+n} = M_{k+n} (e)$$

Define  $r_1(x)$  to be the element represented by:

$$S^{2k} \wedge X^+ \xrightarrow{f} M_{k+n} \subset M_{k+n} (e)$$

with trivial  $G$ -action on  $S^{2k} \wedge X^+$ .

If we forget about the  $G$ -action on  $Y_{k+n}^G$ , we receive a  $(k+n)$  - vector bundle. Given a  $G$ -module  $V \subset V^\infty =$  the universal  $G$ -module of  $G$ , map the standard bundle on  $G_{k+n}(V)$  (after forgetting the action of  $G$ ) isomorphically onto the standard bundle on  $G_{k+n}(C^{1V1}) \subset B_{k+n} = B_{k+n}(e)$ .

This defines a bundle map:

$$E_{k+n}(G) \longrightarrow E_{k+n}$$

and thus induces a map:

$$M_{k+n}(G) \xrightarrow{\nabla} M_{k+n}. \text{ In terms of this map:}$$

$r(r_1(x))$  is represented by:

$$S^{2k} \wedge X^+ \xrightarrow{f} M_{k+n} \subset M_{k+n}(G) \xrightarrow{\nabla} M_{k+n}$$



By the definition of  $\nabla$ , we can assume that the composition:

$$M_{k+n} \subset M_{k+n}(G) \xrightarrow{\nabla} M_{k+n}$$

is  $= \text{id} : M_{k+n} \rightarrow M_{k+n}$ . Hence  $\text{ror}_1 = \text{id}$ . Q.E.D.

§ 1.4 : (B) The Thom Isomorphism Theorem For  $U_G^*(-)$  :-

(i) Let  $\pi : E \rightarrow X$  be an  $n$ -dimensional  $G$ -vector bundle over the compact  $G$ -space  $X$ . By Proposition 1.3.1, we can assign to  $E$  a  $G$ -homotopy class of maps  $[\epsilon : E \rightarrow \text{En}(G)]$ , namely the classifying one. This induces a based  $G$ -homotopy class of maps  $[\epsilon : E^+ \rightarrow \text{Mn}(G)]$ . Denote by  $t_E$  the element of  $U_G^{2n}(E)$  represented by  $[\epsilon : E^+ \rightarrow \text{Mn}(G)]$  and call it the Thom class of  $E$  over  $X$

(ii) Suppose  $\pi' : F \rightarrow X$  is an  $m$ -dimensional  $G$ -vector bundle over  $X$ . We can regard  $E \oplus F$  as a  $G$ -vector bundle over  $E$  in the natural way. Let  $E \oplus F \xrightarrow{P_2} F$  be the natural bundle map,

$$\begin{array}{ccc} E \oplus F & \xrightarrow{P_2} & F \\ P_1 \downarrow & \searrow \pi & \downarrow \pi' \\ E & \xrightarrow{\quad} & X \end{array}$$

$P_1, P_2$  are the projections onto the first and second factors.

$P_2$  induces a  $G$ -map of Thom spaces :  $(E \oplus F)^+ \xrightarrow{P_2} F^+$ . Let

$\epsilon' : F^+ \rightarrow \text{Mn}(G)$  be a representative of the Thom class of

$F$  over  $X$ . Define the Thom class of  $E \oplus F$  over  $F$  to be the element represented by the composition:

$$(E \oplus F)^+ \xrightarrow{P_2} F^+ \xrightarrow{\epsilon'} \text{Mn}(G).$$

(iii)  $E$  is trivial i.e.  $E = V \times X$  for some  $G$ -module  $V$  and  $X$  locally compact. Again we can define the Thom class  $t_E$  to be the one represented by the natural map:

$$(V \times X)^+ \longrightarrow M_{|V|} (G).$$

In each of the situations (i), (ii) or (iii) we can define a Thom homomorphism. Since it is the same kind of construction in (i), (ii) and (iii), we shall do it only for case (i). The bundle map:

$$\begin{array}{ccc} E & \xrightarrow{\pi_X} & X \times E \\ \pi \downarrow & & \downarrow 1 \times \pi \\ X & \xrightarrow{\Delta} & X \times X \end{array}$$

(where  $\Delta$  is the diagonal map) induces a  $G$ -map of Thom spaces  $E^+ \xrightarrow{(\pi \wedge 1)} X^+ \wedge E^+$ . Let  $x \in U_G^{2k}(X)$  be represented by:

$$V^+ \wedge X^+ \xrightarrow{f} M_{|V|+k}(G)$$

Define  $\varphi(x) \in \tilde{U}_G^{2k+2n}(E^+)$  to be the element represented by the composition.

$$V^+ \wedge E^+ \xrightarrow{1 \wedge (\pi \wedge 1)} V^+ \wedge (X^+ \wedge E^+) = (V^+ \wedge X^+) \wedge E^+ \xrightarrow{f \wedge e} M_{|V|+k}(G) \wedge M_n(G) \xrightarrow{(1 \wedge 1)} M_{|V|+k+n}(G)$$

Similarly  $\varphi(x)$  can be defined for  $x \in U_G^{2k-1}(X)$ . The map:

$$\varphi : U_G^*(X) \longrightarrow \tilde{U}_G^*(E^+)$$

is a homomorphism which we call the Thom homomorphism.

Proposition 1.4.4 (tom Dieck [11] )

The Thom homomorphism is transitive i.e. let  $E$  be an  $m$ -dim  $\mathbb{L}$ - $G$ -vector bundle over the compact  $G$ -space  $X$ ,  $F$  an  $n$ -dim  $\mathbb{L}$ - $G$ -vector bundle over  $X$ , then the diagrams:

$$\begin{array}{ccc} U_G^*(X) & \xrightarrow{\varphi} & U_G^*(E \oplus F) \\ \varphi \searrow & & \nearrow \varphi \\ & U_G^*(E) & \end{array} \quad \& \quad \begin{array}{ccc} U_G^*(E) & \xrightarrow{\varphi} & U_G^*(E \oplus E \oplus F) \\ \varphi \searrow & & \nearrow \varphi \\ & U_G^*(E \oplus F) & \end{array}$$

are commutative.

Th. 1.4.5: The Thom Isomorphism Theorem (T. tom Dieck) [11] P.21 )

Given an  $n$ -dim  $\mathbb{L}$ - $G$ -vector bundle  $E$  over a compact  $G$ -space  $X$ , then the Thom homomorphism

$$U_G^k(X) \longrightarrow \tilde{U}_G^{k+2n}(E^+)$$

is an isomorphism. If  $E$  is trivial, it is enough to have  $X$  locally compact.

Proof:

Case 1: The trivial case  $E = V \times X$  :

The classifying map of this bundle is covered by the map

$$V \times X \longrightarrow E_n(G)$$

sending  $(v, x) \longrightarrow (V, v) \quad \forall (v, x) \in V \times X$ . Hence given  $x \in U_G^*(X)$  represented by:

$$W^+ \wedge X^+ \xrightarrow{f} M_{lWl+k}(G)$$

one can check that  $\phi(x)$  is represented by the composition:

$$W^+ \wedge (V \times X)^+ = V^+ \wedge (W^+ \wedge X^+) \xrightarrow{1 \wedge f} V^+ \wedge M_{1W1+k}(G) \downarrow M_{1V1+1W1+k}(G)$$

where the last map comes from the definition of the Thom spectrum. Therefore, by the definition of  $U_G^*(-)$ ,  $\phi$  is a monomorphism. To see that it is an epimorphism, suppose  $y \in \tilde{U}_G^k(E^+)$  is represented by:

$$W^+ \wedge E^+ = W^+ \wedge (V \times X)^+ \xrightarrow{g'} M_{1W1+l}(G)$$

Then  $y = \phi(x)$  where  $x$  is represented by:

$$(W \oplus V)^+ \wedge X^+ = W^+ \wedge (V^+ \wedge X^+) \cong W^+ \wedge (V \times X)^+ \xrightarrow{f'} M_{1W1+l}(G)$$

Case 2 : the general case:

By (Segal [20] Proposition 2.4) ,  $\exists$  a  $G$ -vector bundle  $F$  over  $X$  st.  $E \oplus F$  is trivial, say  $E \oplus F = \underline{V} = V \times X$ .

By proposition 1.4.4 and Case 1 above, we have a commutative

$$\begin{array}{ccc} \text{diagram } (*) : U_G^k(X) & \xrightarrow{\phi} & \tilde{U}_G^{k+21V1}((E \oplus F)^+) \\ & \searrow \phi & \nearrow \phi \\ & \tilde{U}_G^{k+2n}(E^+) & \end{array} \dots\dots\dots(*)$$

in which the horizontal arrow is an isomorphism. Hence

$$\phi : \tilde{U}_G^{k+2n}(E^+) \longrightarrow \tilde{U}_G^{k+21V1}((E \oplus F)^+)$$

is an epimorphism. By the same reasoning, we have a commutative diagram :

$$\begin{array}{ccc}
 \tilde{U}_G^{k+2n}(E^+) & \xrightarrow{\phi} & \tilde{U}_G^{k+2n+21V1}((E \oplus V)^+) \\
 & \searrow \phi & \nearrow \phi \\
 & \tilde{U}_G^{k+21V1}((E \oplus F)^+) &
 \end{array}$$

and the horizontal arrow is an isomorphism. Therefore,  $\phi :$   
 $\tilde{U}_G^{k+2n}(E^+) \longrightarrow \tilde{U}_G^{k+21V1}((E \oplus F)^+)$   
 is a monomorphism. Hence it is an isomorphism. By considering  
 the diagram (\*), the result follows. Q.E.D.

1.4: (C) The Isomorphism  $F(G,H)$  :

Let  $H$  be a closed subgroup of  $G$  and let  $X$  be a compact  
 $H$ -space. Recall that the  $G$ -space  $G \times_H X$  is the space obtained  
 from the cartesian product  $G \times X$  by identifying  $(g,y)$  with  
 $(gh, h^{-1}y) \forall h \in H$  with  $G$  acting on it via:

$$g_1 [g,y] = [g_1 g, y] .$$

Note:

Suppose  $Y$  is a compact  $G$ -space. Then we have a natural  
 identification of  $G$ -spaces:

$$v : G \times_H (Y \times X) \xrightarrow{\sim} Y \times_H (G \times X) \quad (1.4.6)$$

( $Y$  on the left is regarded as an  $H$ -space by restriction) :  
 it sends  $[g, (y,x)] \longrightarrow (gy, [g,x])$ , and thus its inverse  
 is given by :

$$(y, [g,x]) \longrightarrow [g, (g^{-1}y, x)] .$$

There is a canonical homomorphism (T. tom Dieck [11] P.24):

$$F(G,H) : U_G^{**} (G \times_H X) \longrightarrow U_H^* (X)$$

defined as the composition:

$$U_G^* (G \times_H X) \cong \tilde{U}_G^* (G^+ \hat{\wedge}_H X^+) \xrightarrow{r} \tilde{U}_H^* (G^+ \hat{\wedge}_H X^+) \xrightarrow{q^*} \tilde{U}_H^* (X^+) = U_H^* (X)$$

where  $q^*$  is induced by the map:

$$q : X^+ \longrightarrow G^+ \hat{\wedge}_H X^+$$

defined by :  $x \rightarrow [1, x]$ . tom Dieck ([11] P.24) has shown too that  $F(G,H)$  is an isomorphism with invese:

$$F(H,G) : U_H^* (X) \longrightarrow U_G^* (G \times_H X) .$$

$F(H,G)$  is defined by the composition:-

$$\begin{aligned} & [V^+ \wedge X^+ , M_k(H)]_H^{\circ} \xrightarrow{(1)} \\ & [G^+ \hat{\wedge}_H (V^+ \wedge X^+) , G^+ \hat{\wedge}_H M_k(H)]_G^{\circ} \xrightarrow{(2)} \\ & [G^+ \hat{\wedge}_H (V^+ \wedge X^+) , M_k(G)]_G^{\circ} \xrightarrow{(3)} \\ & \tilde{U}_G^* (G^+ \hat{\wedge}_H (V^+ \wedge X^+) ) \stackrel{(4)}{\cong} U_G^* (G \times_H X) \end{aligned}$$

where:

$$(1) \text{ sends } (f: V^+ \wedge X^+ \longrightarrow M_k(H)) \xrightarrow{t_0} 1 \wedge f : G^+ \hat{\wedge}_H (V^+ \wedge X^+) \longrightarrow G^+ \hat{\wedge}_H M_k(H),$$

$$(2) \text{ is induced by the bundle map: } G \times_H E'_k(H) \longrightarrow E_k(G),$$

(3) comes from the definition of  $U_G^*$ , & (4) is the invese of the Thom isomorphism (Th 1.4.2) applied to the  $G$ -vector bundle  $G \times_H (V \times X) \longrightarrow G \times_H X$ .

Lemma 1.4.7:

Let  $X$  be a compact  $G$ -space, and  $H$  a closed subgroup of  $G$ . Let  $P_2 : G/H \times X \rightarrow X$  be projection onto the 2nd factor. Then we have a commutative diagram:

$$\begin{array}{ccc}
 U_G^*(X) & \xrightarrow{P_2^*} & U_G^*(G/H \times X) \\
 \downarrow r & & \downarrow v^* \cong \\
 U_H^*(X) & \xrightarrow{F(H,G)} & U_G^*(G \times_H X)
 \end{array}$$

where the isomorphism  $U_G^*(G/H \times X) \cong U_G^*(G \times_H X)$

is the one induced by the identification (1.4.6):

$$v : G \times_H X = G/H \times X,$$

Proof:

Since  $F(H,G) : U_H^*(X) \rightarrow U_G^*(G \times_H X)$  is an isomorphism with inverse  $F(G,H) : U_G^*(G \times_H X) \rightarrow U_H^*(X)$ , it is enough to prove that the diagram :

$$\begin{array}{ccc}
 U_G^*(X) & \xrightarrow{P_2^*} & U_G^*(G/H \times X) \\
 \downarrow r & & \downarrow v^* \cong \\
 U_H^*(X) & \xleftarrow{F(G,H)} & U_G^*(G \times_H X)
 \end{array} \quad \dots\dots\dots (d_2)$$

is commutative.

Suppose  $x \in U_G^*(X)$  is represented by

$V^+ \wedge X^+ \xrightarrow{f} M_k(G)$ . Then  $P_2^*(x)$  is represented by the composition:

$V^+ \wedge (G/H \times X)^+ \xrightarrow{1 \wedge P_2} V^+ \wedge X^+ \xrightarrow{f} M_k(G)$ .  $v^*(P_2^*(x))$  is represented by the composition  $S_1$ :

$$V^+ \wedge (G \times_H X)^+ \xrightarrow{1 \wedge v} V^+ \wedge (G/H \times X)^+ \xrightarrow{1 \wedge P_2} V^+ \wedge X^+ \xrightarrow{f} M_k(G)$$

$S_1$  is a  $G$ -map. Denote it by  $S_1'$  when regarded as an  $H$ -map by restriction.

$F(G,H)(v^*(P_2^*(x)))$  is represented by the composition:

$$V^+ \wedge X^+ \xrightarrow{1 \wedge q} V^+ \wedge (G \times_H X)^+ \xrightarrow{S_1'} M_k(G) \xrightarrow{\alpha} M_k(H)$$

where  $\alpha$  is obtained by regarding  $V_k^G$  as an  $H$ -vector bundle by restriction. Now the  $H$ -map given by the composition:

$$V^+ \wedge X^+ \xrightarrow{1 \wedge q} V^+ \wedge (G \times_H X)^+ \xrightarrow{1 \wedge v} V^+ \wedge (G/H \times X)^+ \xrightarrow{1 \wedge P_2} V^+ \wedge X^+$$

$$= V^+ \wedge X^+ \xrightarrow{\text{id} \wedge \text{id}} V^+ \wedge X^+ . \text{ Therefore,}$$

$$F(G,H)(v^*(P_2^*(x))) = r(x) \text{ i.e. the diagram } (d_2)$$

is commutative.

Q.E.D.

### § 1.5 (a): Spectral Sequences:-

Our aim is to set up the  $U_G^*$  - spectral sequence, and give an application of it in § 1.5.b. We assume a number of definitions all of which can be found in (Segal [20][21]).



Let  $h_G^* = (h_G^k)$  be an equivariant cohomology theory on the category  $(\underline{G}_1)$  (see 1.1). Let  $F = (F_\alpha)_{\alpha \in A}$  be a closed finite covering of a locally compact  $G$ -space  $X$ . Let  $N_F$  be the nerve of  $F$  - a finite simplicial complex. Let  $|N_F|$  be its geometrical realization. Define  $W_F = \bigcup_{\sigma \in \Sigma} (F \times |\sigma|)$  where  $\sigma$  runs through the finite subsets of  $A$  s.t.  $\bigcap_{\alpha \in \sigma} F_\alpha \neq \emptyset$ .  $W_F$  is a closed  $G$ -subspace of  $X \times |N_F|$ . Define

$$w : W_F \rightarrow X$$

to be projection onto the first factor - a proper  $G$ -map since  $|N_F|$  is compact. Now define a filtration of  $W_F$  by:

$$W_F^P = \bigcup_{\dim \sigma \leq P} (F \times |\sigma|).$$

Proposition 1.5.1:

The natural homomorphism:

$$w^* : h_G^*(X) \rightarrow h_G^*(W_F)$$

is an isomorphism.

Proof:

By the method of Segal (Atiyah-Segal [4] Proposition 3.1.3). In the manner of Cartan-Eilenberg ([10] P.333), we associate to the filtration:

$$W_F \supset \dots \supset W_F^P \supset \dots \supset W_F^0$$

a spectral sequence whose  $E_2^{p,q}$  term is the  $p$ th cohomology of the complex

$$h_G^q(W^0) \rightarrow h_G^{q+1}(W_F^1, W_F^0) \rightarrow \dots \rightarrow h_G^{p+q}(W_F^P, W_F^{P-1}) \rightarrow \dots$$

& with termination  $h_G^*(W_F) \cong h_G^*(X)$  (Proposition 1.5.1)

By a standard argument (Segal [4]§3.2) [21] Proposition 5.1) one can verify

$$(i) \quad h_G^{P+q}(W_F^P, W_F^{P-1}) = \prod_{\dim \sigma = p} h_G^q(F \sigma) \text{ i.e. } E_1^{P,q} = \prod_{\substack{G \\ \dim \sigma = P}} h_G^q(F \sigma)$$

$$(ii) \text{ \& the differential } d : E_1^{P,q} \longrightarrow E_1^{P+1,q}$$

corresponds to the differential of the complex of cochains of  $N_F$  with coefficients in the system :  $\sigma \rightarrow h_G^q(F \sigma)$ . So:

Proposition 1.5.2:

Let  $F$  be a finite closed covering of a locally compact  $G$ -space  $X$ . There is a spectral sequence  $H^P(N_F; h_G^q(F)) \Rightarrow h_G^*(X)$  where  $h_G^q(F)$  means the coefficient system  $\sigma \rightarrow h_G^q(F \sigma)$ .

Proposition 1.5.3:

Let  $X$  be a locally compact  $G$ -space,  $Y$  a compact  $G$ -space on which  $G$  acts trivially and  $f: X \rightarrow Y$  a  $G$ -map. There is a spectral sequence  $H^P(Y; h_G^q f) \Rightarrow h_G^*(X)$  where  $h_G^q f$  is the sheaf on  $Y$  associated to the presheaf :  $V \rightarrow h_G^q(f^{-1}(\bar{V}))$ . If in addition  $h_G^*$  is continuous (see Proposition 1.2.2),  $h_G^q f$  has stalk  $h_G^q(f^{-1}y)$  at the point  $y$ .

Proof:

By the method of Segal [21] Proposition 5.2. If  $F$  is a finite open covering of  $Y$ , form the spectral sequence  $E(F)$  for the finite closed covering  $f^{-1}\bar{F}$  of  $X$  (see Proposition 1.5.2). This terminates with  $h_G^*(X)$  and begins with the Čech cohomology of the covering  $F$  with coeffs in the presheaf

$V \rightarrow h_G^q (f^{-1} \bar{V})$ . Since  $Y$  is compact, the set  $S$  of finite open coverings of  $Y$  is cofinal in the set of open coverings of  $Y$ . Therefore, if we take  $\varinjlim \{E(F)\}$ , and use the fact that the Čech cohomology of a compact space with coeffs in a presheaf is = its cohomology with coeffs in the associated sheaf (Spanier [22] Chap 6), the result follows. Q.E.D. (for more details see Segal [21]). By Cor. 1.3.4 :

Cor 1.5.4:

Proposition 1.5.3 is true when  $h_G^* = U_G^*$ . In particular, let  $X$  be a compact  $G$ -space,  $Y = X/G$  &  $\pi: X \rightarrow X/G$  be projection. There is a spectral sequence  $H^p(X/G; \mathcal{U}_G^q) \Rightarrow U_G^*(X)$  where  $\mathcal{U}_G^q$  is the sheaf on  $X/G$  associated to the presheaf  $V \rightarrow U_G^q(\pi^{-1} \bar{V})$ . The stalk of  $\mathcal{U}_G^q$  at an orbit  $Gx = G/G_x$  is  $U_G^q(G/G_x)$  where  $G_x$  is the stabilizer of  $x$ .

§ 1.5 (b) : Free Group Action:

Let  $G, K$  be compact Lie groups, and  $X$  a compact  $(G \times K)$ -space s.t.  $K$  acts freely on  $X$ . Let  $pr: X \rightarrow X/K$  be the projection. It induces a homomorphism :

$$pr^* : U_G^*(X/K) \longrightarrow U_{G \times K}^*(X)$$

defined by the composition:

$$[V^+ \wedge (X/K)^+, M_n(G)]_G^0 \xrightarrow{(1)}$$

$$[V^+ \wedge X^+, M_n(G)]_G^0 \xrightarrow{(2)}$$

$$[V^+ \wedge X^+, M_n(G \times K)]_{G \times K}^0 \xrightarrow{(3)} U_{G \times K}^*(X),$$

where (1) is induced by  $\text{pr} : X \rightarrow X/K$ , (2) is induced by the bundle map  $E_n(G) \rightarrow E_n(G \times K)$  when  $E_n(G)$  is regarded as a  $G \times K$ -bundle by giving it trivial  $K$ -action, and (3) is the natural map.  $V$  is regarded as a  $G \times K$ -module as follows:

$$(g,k)v = gv \quad \forall v \in V, (g,k) \in G \times K.$$

Proposition 1.5.5:

Suppose  $G \times K$  acts on a compact space  $X$  s.t. the action of  $K$  on  $X$  is free. Then :

$$\text{pr}^* : U_G^*(X/K) \rightarrow U_{G \times K}^*(X)$$

is an isomorphism.

Proof:

Let  $\{ {}_1E_r \}$  be the spectral sequence  $H^P((X/K)/G; \mathcal{U}_G^q) \Rightarrow U_G^*(X/K)$  where  $\mathcal{U}_G^q$  is the sheaf on  $(X/K)/G$  associated to the presheaf  $V \rightarrow U_G^q(\pi^{-1}\bar{V})$  and  $\pi : X/K \rightarrow (X/K)/G$  is projection and let  $\{ {}_2E_s \}$  be the spectral sequence :  $H^P(X/G \times K; \mathcal{U}_{G \times K}^q)$  where  $\mathcal{U}_{G \times K}^q$  is the sheaf on  $X/G \times K = (X/K)/G$  associated to the presheaf  $V \rightarrow U_{G \times K}^q(\pi_1^{-1}\bar{V})$  ( $\pi_1 : X \rightarrow X/G \times K$  is projection) (see Cor 1.5.4). Now  $\text{pr}^* : U_G^*(X/K) \rightarrow U_{G \times K}^*(X)$  induces a homomorphism of spectral sequences :  $\{ {}_1E_r \} \rightarrow \{ {}_2E_r \}$  in the natural way. If we can show that this is an isomorphism at the  $E_2$ -level, it would follow that it is an isomorphism at each  $E_r$ -level  $r \geq 2$  (Cartan - E. Jenberg [10])

and hence the result would follow.

By Cor 1.5.4, the stalk of  $\mathcal{U}_G^q$  at an orbit  $G[x]$  is  $U_G^q(G/G')$  where  $G'$  is the stabilizer of  $[x] \in X/K$ , and the stalk of  $\mathcal{U}_{GxK}^q$  at the corresponding orbit  $(GxK)x$  is  $U_{GxK}^q((GxK)/G'')$  where  $G'' \subset GxK$  is the stabilizer of  $x \in X$ . The homomorphism :

$$h : G'' \longrightarrow G'$$

given by projection onto the first factor is an isomorphism.

For it is clearly epi. To see that it is mono, suppose

$(g, k_1) \cdot x = x = (g, k_2) \cdot x$ . Since  $K$  acts freely on  $X$ , then

$k_1 = k_2$  and so  $G'' \xrightarrow{h} G'$  is an isomorphism. We can also

identify  $((GxK)/G'')/K$  and  $G/G'$  by the projection map.

Now the diagram :

$$\begin{array}{ccc} U_G^q(G/G') & \xrightarrow{\text{Pr}^*} & U_{GxK}^q((GxK)/G'') \\ \downarrow F(G, G') & & \downarrow F(GxK, G'') \\ U_{G'}^q & \xrightarrow[\text{Rh}]{\approx} & U_{G''}^q \end{array} \quad \text{by Proposition 1.4.1}$$

is commutative and the two vertical arrows are isomorphisms

(§ 1.4B). Hence  $U_G^q(G/G') \xrightarrow{\text{Pr}^*} U_{GxK}^q((GxK)/G'')$  is

an isomorphism.

Q.E.D.

## § 1.6 : Multiplicative Equivariant Cohomology Theories:-

As we have demonstrated in the previous section, one of the uses of Proposition 1.5.3 is to enable us to reduce the solution of a number of problems to investigating what happens when only orbits

are involved. The proposition we are aiming for now plays a similar role when dealing with multiplicative theories.

Definition 1.6.1:

A multiplicative equivariant cohomology theory on  $(\underline{c}_1)$  is an equivariant cohomology theory,  $h_G^*$ , on  $(\underline{c}_1)$  such that (i) for each  $(X, A)$ ,  $(X^1, A^1)$  in  $(\underline{c}_1)$ , there is a homomorphism:

$$\otimes : h_G^i(X, A) \otimes h_G^j(X^1, A^1) \longrightarrow h_G^{i+j}(X \times X^1, X \times A^1 \cup A \times X^1)$$

which is associative, anticommutative, natural under proper  $G$ -maps of pairs in  $(\underline{c}_1)$ , has a unit  $1 \in h_G^0(\text{pt})$ ,

(ii) Let  $X = X^1$  & let :

$$h_G^i(X, A) \otimes h_G^j(X, A^1) \xrightarrow{U} h_G^{i+j}(X, A \cup A^1)$$

be the induced internal pairing i.e. the composition:

$$\begin{aligned} h_G^i(X, A) \otimes h_G^j(X, A^1) &\xrightarrow{\otimes} h_G^{i+j}(X \times X, X \times A^1 \cup A \times X) \\ &\xrightarrow{\Delta^*} h_G^{i+j}(X, A \cup A^1) \end{aligned}$$

where  $\Delta : X \longrightarrow X \times X$  is the diagonal map. We require that under this internal pairing,  $h_G^*(A) \xrightarrow{\delta} h_G^*(X, A)$  is an  $h_G^*(X)$  - module homomorphism.

As an example, we have

Lemma 1.6.2:

$U_G^*$  is a multiplicative equivariant cohomology theory.

Proof:-

Let  $X, X^1$  be compact  $G$ -spaces with base point,  $A, A^1$  closed  $G$ -subspaces of  $X, X^1$  respectively, and containing the base points.

Let  $x \in \tilde{U}_G^{2i}(X)$ ,  $x^1 \in \tilde{U}_G^j(X^1)$  be represented by:

$f: V^+ \wedge X \rightarrow M_{1V1+i}(G)$ , and  $f': W^+ \wedge X' \rightarrow M_{1W1+j}(G)$ .

Define  $x \otimes x' \in \tilde{U}_G^{2i+2j}(X \wedge X')$  to be the element represented by:

$$(V \oplus W)^+ \wedge X \wedge X^1 = (V^+ \wedge X) \wedge (W^+ \wedge X^1) \xrightarrow{f \wedge f'} M_{1V1+i}(G) \wedge M_{1W1+j}(G) \\ \xrightarrow{(1V1+i, 1W1+j)} M_{1V \oplus W1+i+j}(G)$$

This defines a multiplication:

$$\otimes : \tilde{U}_G^*(XUCA) \otimes \tilde{U}_G^*(X^1 UCA) \rightarrow \tilde{U}_G^*((XUCA) \wedge (X^1 UCA^1)) \cong \\ \cong \tilde{U}_G^*(X \wedge X^1 UC (X \wedge X^1 U A \wedge X^1))$$

where the last isomorphism is induced by a natural proper  $G$ -homotopy equivalence. Hence given  $(X, A)$ ,  $(X', A') \in (C_1)$ , we receive a homomorphism:

$$\otimes : U_G^*(X, A) \otimes U_G^*(X', A') \rightarrow U_G^*(X \times X'; A \times X' U X \times A').$$

It is easy to see that this is associative, anticommutative, natural under morphisms in  $(C_1)$ . In § 1.4, we defined

$1 \in U_G^0(\text{point})$ . It can also be checked that (ii) holds.

Q.E.D.

### Proposition 1.6.3:

Let  $h_G^*$  be a multiplicative equivariant cohomology theory.

Let  $\pi : B \rightarrow X$  be a  $G$ -map of compact  $G$ -spaces and  $\gamma_1, \dots, \gamma_n$  homogeneous elements of  $h_G^*(B)$ . Let  $N^*$  be the free  $h_G^*$ -module

generated by  $\gamma_1, \dots, \gamma_n$  ( $h_G^* = h_G^*$  (point)). Suppose every orbit in  $X$  has a closed  $G$ -neighbourhood  $F$  s.t.  $\forall$  closed  $G$ -subspace  $F_1$  of  $F$ , the natural map:

$$h_G^*(F_1) \otimes_{h_G^*} N^* \longrightarrow h_G^*(\pi^{-1} F_1)$$

is an isomorphism. Then for any closed  $G$ -subspace  $Y$  of  $X$ , the map:

$$h_G^*(X, Y) \otimes_{h_G^*} N^* \longrightarrow h_G^*(B, \pi^{-1}(Y))$$

is an isomorphism.

Proof:

By the method of proof of Th. 2.7.8, Atiyah [1].

Q.E.D.



## Chapter 2: The Equivariant Cobordism Theory of Projective Spaces:

### § 2.1 (a) The Conner-Floyd Map $\mu$ :

2.1(a): Given a locally compact  $G$ -space  $X$  and a closed  $G$ -subspace  $A$  of  $X$ , let  $K_G^*(X, A)$  be defined as in Segal [22]. We shall assume the following result:

#### Th 2.1.1 :

Let  $E$  be an  $n$ -dimensional  $G$ -vector bundle over a locally compact  $G$ -space  $X$ . Then the Thom homomorphism (Segal [20])

$$q_* : K_G^*(X) \longrightarrow K_G^*(E)$$

is an isomorphism.

In contrast to the Thom isomorphism theorem for  $U_G^*$ , the proof of which is immediate from the definition and the transitivity of the Thom homomorphism (Th 1.4.5), Th 2.1.1 had to be proved (for  $E$  trivial) by using elliptic operators (Atiyah [2]). By means of Th 2.1.1, we would like to define the Conner-Floyd map:

$$\mu : U_G^*( - ) \longrightarrow K_G^*( - )$$

(see C-F [8] P.28 for the case  $G=e$ ). Given a  $G$ -module  $V$ , let

$M_k(V)$  denote the Thom space of the standard bundle  $E_k(V)$  over

$G_k(V)$  = the Grassmann Manifold (of  $k$ -subspaces) of  $V$ . If

$V \leq W$  (i.e.  $\exists V'$  s.t.  $V \oplus V' = W$ ), then inclusion  $E_k(V)$

$\hookrightarrow E_k(W)$  induces a  $G$ -map  $M_k(V) \longrightarrow M_k(W)$  and hence a homomorphism:

$$a_V^W : \tilde{K}_G(M_k(W)) \longrightarrow \tilde{K}_G(M_k(V))$$

Define  $\tilde{O}(M_k(G)) = \varprojlim \{M_k(V)\}; a_v^W\}$ .

Let  $\tilde{\lambda}_{E_k}(V)$  be the Thom class in  $\tilde{K}_G(M_k(V))$  ([20] P.140).

Then  $a_v^W(\tilde{\lambda}_{E_k}(W)) = \tilde{\lambda}_{E_k}(V)$  This defines a natural element  $\tilde{\lambda}^k$  of  $\tilde{O}(M_k(G))$ .

Definition:

Let  $X$  be a locally compact  $G$ -space. Suppose  $x \in \tilde{U}_G^{2n}(X^+)$  is represented by  $f: V^+ \wedge X^+ \longrightarrow M_{1V1+n}(G)$ . By compactness of  $V^+ \wedge X^+$ , we can find a  $G$ -module  $W$  s.t.  $f(V^+ \wedge X^+) \subset M_{1V1+n}(W)$ . Define  $\mu(x) \in K(X)$  to be the image of  $\tilde{\lambda}^L$  under the

composition:  
 $(b_1) \dots \tilde{O}(M_L(G)) \xrightarrow{(1)} K_G(M_L(W)) \xrightarrow{f^*} \tilde{K}_G(V^+ \wedge X^+) \xrightarrow{(2)} K_G(X)$   
 where  $L = 1V1+n$ , (1) comes from the definition of  $\tilde{O}(M_L(G))$ , and (2) is the inverse of the Thom isomorphism for the trivial bundle  $V \times X \longrightarrow X$  (Atiyah [2]).

Lemma 2.1.1 :

$$\mu : U_G^*(X) \longrightarrow K_G^*(X)$$

is well-defined:

Proof:

(i) To prove  $\mu$  does not depend on the choice of  $W$ :

Suppose  $W_1$  and  $W_2$  are s.t.  $f(V^+ \wedge X^+) \subset M_L(W_i)$ ,  $i = 1, 2$ . Hence  $f(V^+ \wedge X^+) \subset M_L(W_1 \oplus W_2)$ , and for  $i = 1, 2$ , we have a commutative diagram:

$$\begin{array}{ccc} \tilde{O}(M_L(G)) & \xrightarrow{(1)} & \tilde{K}_G(M_L(W_i)) \\ (1) \downarrow & & \downarrow f^* \\ \tilde{K}_G(M_L(W_1 \oplus W_2)) & \xrightarrow{f^*} & \tilde{K}_G(V^+ \wedge X^+) \end{array}$$

where the last map is that of the Thom spectrum. Put

$\iota_1 = 1V_11$ . Using this representative of  $x$ ,  $\mu(x)$  is the

image of  $\overline{\lambda}^{\iota_1 + \iota_1}$  in the composition:

$$(b_2) \dots \tilde{O}(M_{\iota_1 + \iota_1}(G)) \xrightarrow{(1)} \tilde{K}_G(M_{\iota_1 + \iota_1}(V_1 \oplus W)) \xrightarrow{f_1^*} \tilde{K}_G((V_1 \oplus V)^+ \wedge X^+) \cong \tilde{K}_G(X).$$

By the fact that  $\lambda_{E \otimes F} = \lambda_E \cdot \lambda_F$  (Segal [20]), and the multiplicativity of the maps (1),  $f^*$  and  $f_1^*$ , it follows that the composition  $(b_1)$  = the composition  $(b_2)$ .

Q.E.D.

Lemma 2.1.2:

$$\mu : U_G^*(-) \longrightarrow K_G^*(-)$$

is a natural multiplicative transformation of the two theories:

Proof:

$\mu$  is multiplicative because the Thom class (when it  $\exists$ ) in

$K_G$ -theory satisfies  $\lambda_{E \otimes F} = \lambda_E \cdot \lambda_F$  (Segal [20]).

The rest of the statement follows from the definition of  $\mu$ .

Q.E.D.

2.1(b): The Thom class of a line bundle in  $K_G$ -theory:

Let  $L$  be a 1-dim  $\mathbb{C}$ -G-vector bundle over a compact G-space  $X$ .

Let  $p(L)$  be the associated principal  $U(1)$ -bundle. Recall that

the join  $p(L) \circ U(1)$  consists of all points of the form

$(1-t)e + tu$  for  $0 \leq t \leq 1$ ,  $e \in p(L)$ ,  $u \in U(1)$ .  $U(1)$  acts

principally on it via:

$$((1-t)e + tu)v = (1-t)ev + t uv \text{ and } G \text{ acts on it as}$$

$$\text{follows: } ((1-t)e + tu)g = (1-t)ge + tu.$$

Lemma 2.1.3:  $\exists$  a canonical  $G$ -homeomorphism:

$$f: L^+ \longrightarrow p(L) \circ U(1) / U(1) .$$

Proof:

As in the ordinary case,  $G=\mathbb{C}$  (Conner - F [8] P.19 ).

We identify the disc bundle of  $L$ ,  $D(L)$  with  $p(L) \times_{U(1)} D^2$  and

$S(L)$  = the sphere bundle of  $L$  with  $p(L) \times_{U(1)} S^1$

where  $D^2$  and  $S^1$  are the unit disc and unit sphere in  $\mathbb{C}$  =

the complex plane. Define

$$f' : p(L) \times D^2 \longrightarrow p(L) \circ U(1)$$

$$\text{by : } (e, d) \longrightarrow (1 - |d|^2) e + |d|^2 ( \bar{d} / |d| ) .$$

$f'$  is equivariant w.r.t.  $U(1)$  - actions. So there is defined

a  $G$ -map :

$$f : (D(L) , S(L)) \longrightarrow (p(L) \circ U(1)) /_{U(1)} , \quad x_0$$

where  $x_0$  is the orbit containing all  $1 \cdot u, u \in U(1)$ .

This induces a  $G$ -map:

$$D(L) /_{S(L)} \longrightarrow (p(L) \circ U(1)) /_{U(1)}$$

which we denote, too, by  $f$ .  $f$  is 1 - 1 and onto and so it is

a homeomorphism since all spaces are compact Hausdorff . Q.E.D.

Proposition 2.1.4:

Let  $L$  be a 1-dim  $\mathbb{C}$ -vector bundle over a compact  $G$ -space  $X$ .

I identify  $X \stackrel{L}{=} \text{the Thom space of } L \text{ with } (p(L) \circ U(1)) /_{U(1)}$

as in 2.1.3. Let  $L'$  be the vector bundle associated to the

principal  $U(1)$  - bundle :  $p(L) \circ U(1)$  over  $X^L$  .

5

Then the Thom class of  $L = 1 - L^*$  in  $\tilde{K}_G(X^L)$ .

Proof:

Observe that the definition of the Thom class of an  $n$ -dimensional  $G$ -vector bundle in  $K_G$ -theory as in Conner-Floyd ([8]) agrees with the definition of Atiyah [1] (P.98-101) because Lemma 2.6.13 in [1] is valid in the equivariant case. Hence proceeding as in C-F [8] Th 4.2, we get the result. Q.E.D.

Cor 2.1.5:

Let  $V$  be a  $G$ -module and  $H$  the Hopf bundle over  $P(V)$ . I identify  $P(V)^H$  with  $P(V \oplus 1)$  as in Lemma 2.1.3 (see also [8] P.21). Then  $\lambda_H = 1 - H_1$  in  $\tilde{K}_G(P(V \oplus 1))$  where  $H_1$  is the Hopf bundle over  $P(V \oplus 1)$ .

Proof:

Because  $S(V \oplus 1) = S(V) \circ U(1)$ ,  $P(V \oplus 1) = S(V \oplus 1)/U(1) = (S(V) \circ U(1))/U(1)$ . Apply Proposition 2.1.4. Q.E.D.

Cor 2.1.6:

The natural element  $\bar{\lambda}^1$  in  $\tilde{O}(M_1(G))$  is represented by  $1 - H \in \tilde{K}_G(P(V \oplus 1)) \quad \forall \quad G\text{-module } V$  where  $H$  is the Hopf bundle over  $P(V \oplus 1)$ .

Lemma 2.1.7:

The Conner-Floyd map  $\mu$  sends the Thom class of a  $G$ -vector bundle in  $U_G^*$ -theory to its Thom class in  $K_G$ -theory (when they are defined).

## § 2.2 : Some Remarks on the Structure of $U_G^*(P(V))$ :-

Suppose  $V$  is a  $G$ -module. Let  $P(V)$  be the projective space of  $V$ , and  $H = \{(L, x) : L \in P(V), x \in L\}$  the Hopf bundle over  $P(V)$ . We can identify the Thom space of  $H$  with  $P(V \oplus 1)$  as indicated in Lemma 2.1.3 and the proof of Cor 2.1.5. So  $M_1(G)$  can be identified with  $P(V^\infty \oplus 1)$

### Definition 2.2.1:

The natural element  $\rho \in U_G^2(P(V))$  is the element represented by:

$$i : P(V)^+ \longrightarrow M_1(G)$$

where  $i : P(V)$  is inclusion. Define  $\rho_H = \rho \in U_G^2(P(V))$ .

This enables us to assign to every  $G$ -line bundle  $L$  over a compact  $G$ -space  $X$  an element  $\rho_L \in U_G^2(X)$  as follows.

As in the proof of Proposition 1.3.1,  $\exists$  a  $G$ -module  $W$  and a  $G$ -map  $f : X \longrightarrow P(W)$  s.t.  $f^*(H_1) = L$  where  $H_1$  is the Hopf bundle over  $P(W)$ . Let  $\rho$  be the natural element in  $U_G^2(P(W))$ . Define  $\rho_L = f^*(\rho)$  where

$$f^* : U_G^*(P(W)) \longrightarrow U_G^*(X)$$

is the induced homomorphism. To see that  $\rho_L$  does not depend on the choice of  $W$  or  $f$ , suppose  $(W', f')$  is used to define  $\rho_L$ . Let  $j : P(W) \longrightarrow P(W \oplus W')$ ,  $j' : P(W') \longrightarrow P(W \oplus W')$  be the inclusion and let  $\rho, \rho', \rho''$  be the natural elements in  $P(W)$ ,  $P(W')$ ,  $P(W \oplus W')$  respectively.

Proposition 2.2.3:

(i)  $U_G^*(P(\mathbb{C}^n))$  is a free  $U_G^*$  - module generated by:

$1, \bar{\rho}_n, \bar{\rho}_n^2, \dots, \bar{\rho}_n^{n-1}$  where  $\bar{\rho}_n \in U_G^2(P(\mathbb{C}^n))$  is the natural element, and subject to the relation:

$$(ii) \bar{\rho}_n^n = 0.$$

Proof:

(ii)  $\bar{\rho}_n = r_1(\rho_n)$  and  $r_1$  is multiplicative. So  $\bar{\rho}_n^n = r_1(\rho_n^n) = 0$ .

(i) By induction. For  $n = 1$ ,  $P(\mathbb{C}^1) = \text{point}$  and the result is obviously true.

Suppose  $U_G^*(P(\mathbb{C}^k))$  is a free  $U_G^*$  - module generated by:

$1, \bar{\rho}_k, \dots, \bar{\rho}_k^{k-1}$ . Let  $i: P(\mathbb{C}^k) \hookrightarrow P(\mathbb{C}^{k+1})$  be the inclusion.

Since  $i^*(\bar{\rho}_{k+1}) = \bar{\rho}_k$ , the cohomology exact sequence of the pair  $(P(\mathbb{C}^{k+1}), P(\mathbb{C}^k))$  gives us a s.e.s.

$$0 \rightarrow U_G^*(P(\mathbb{C}^{k+1}), P(\mathbb{C}^k)) \xrightarrow{\pi^*} U_G^*(P(\mathbb{C}^{k+1})) \xrightarrow{i^*} U_G^*(P(\mathbb{C}^k)) \rightarrow 0$$

It is a split s.e.s. because by hypothesis  $U_G^*(P(\mathbb{C}^k))$  is a free

$U_G^*$  - module. Now  $U_G^*(P(\mathbb{C}^{k+1}), P(\mathbb{C}^k)) \cong \tilde{U}_G^*(S^{2k})$  and so

is a free  $U_G^*$  - module generated by the Thom class,  $\bar{t}$  say, in  $\tilde{U}_G^*(S^{2k})$ .  $-\bar{t} = r_1(t)$  where  $t$  is the Thom class in  $\tilde{U}^*(S^{2k})$

and  $r_1: \tilde{U}^*(S^{2k}) \rightarrow \tilde{U}_G^*(S^{2k})$  is given by L1.4.3.

From (2.2.2), we deduce that  $\rho_{k+1}^k$  and  $\pi^*(t)$  differ by a unit of  $U^*$  i.e.  $\rho_{k+1}^k$  is either  $\pi^*(t)$  or  $-\pi^*(t)$  (Milnor [17]).

Therefore, by naturality,

$$\bar{\rho}_{k+1}^k \text{ is either } \pi^*(t) \text{ or } -\pi^*(t).$$

Thus  $U_G^*(P(\mathbb{C}^{k+1}))$  is freely generated by:

$$1, \bar{\rho}_{k+1}, \bar{\rho}_{k+1}^2, \dots, \bar{\rho}_{k+1}^k$$

The induction is complete.

Q.E.D.

It turns out, however, that  $U_G^*(P(V))$  is not in general a free  $U_G^*$ -module on generators:

$$1, \rho, \dots, \rho^{1V1-1}.$$

T. tom Dieck has a counterexample for  $G = \mathbb{Z}_2$ .

2.3

### Localization:

Let  $\Lambda_G = \{x: x \in U_G^*(\text{point}), \mu(x) = 1 \in RG\}$ . Introduce the elements of  $\Lambda_G$  as denominators in  $U_G^*$  and let the resulting ring be denoted by  $U_G^*[\Lambda_G^{-1}]$ . There is a natural homomorphism  $U_G^* \xrightarrow{(n)} U_G^*[\Lambda_G^{-1}]$  ([6]). This induces a  $U_G^*$ -module structure on  $U_G^*[\Lambda_G^{-1}]$ .

Now for an object  $(x, A)$  in  $(\underline{c}_1)$ , we define:

$$U_G^*(x, A)[\Lambda_G^{-1}] = U_G^*(x, A) \otimes_{U_G^*} U_G^*[\Lambda_G^{-1}]. \text{ Given a morphism}$$

$f$  in  $(\underline{c}_1)$ , define  $U_G^*(-)[\Lambda_G^{-1}](f) = U_G^*(f) \otimes \text{id}$  where  $\text{id}:$

$U_G^*[\Lambda_G^{-1}] \rightarrow U_G^*[\Lambda_G^{-1}]$  is the identity. In what follows, we shall write  $\Lambda$  instead of  $\Lambda_G$  when there is no fear of confusion.

### Proposition 2.3.1:

$U_G^*(-)[\Lambda_G^{-1}]$  is a multiplicative equivariant cohomology theory on the category  $(\underline{c}_1)$ .

### Proof:

The only non-obvious point is exactness of the cohomology sequence. This follows from the fact proved in [6] P.88 that the quotient ring, e.g.  $U_G^*[\Lambda_G^{-1}]$ , is flat when regarded as a module over the ring from which it is constructed. Q.E.D.



What we would like to do is to compute the  $U_G^*(-)[\wedge^{-1}]$  theory of projective spaces,  $P(V)$  ( $V$  a  $G$ -module).

Lemma 2.3.2:

Let  $R$  be a commutative ring with unit 1, and let  $S$  be a subset of  $R$  closed under multiplication. Let  $R[S^{-1}]$  be the quotient ring obtained from  $R$  by taking the elements of  $S$  as denominators and  $R \xrightarrow{(n)} R[S^{-1}]$  the natural map ([6]) Given a ring homomorphism  $h: R \rightarrow R'$  s.t.  $h(s)$  is invertible  $\forall s \in S$ , then  $\exists$  a canonical homomorphism:

$$\bar{h}: R[S^{-1}] \rightarrow R'$$

$$\text{such that the diagram: } \begin{array}{ccc} R & \xrightarrow{(n)} & R[S^{-1}] \\ & \searrow h & \downarrow \bar{h} \\ & & R' \end{array}$$

is commutative.

Proof:

Define  $\bar{h}: R[S^{-1}] \rightarrow R'$  via:  $\bar{h}([x/y]) = h(x)(h(y))^{-1}$ .  
Q.E.D.

Cor 2.3.3:

There a multiplicative natural transformation of cohomology theories:

$$\bar{\mu}: U_G^*(-)[\wedge^{-1}] \rightarrow K_G^*(-)$$

Proof:

Define  $\bar{\mu}: U_G^*[\wedge^{-1}] \rightarrow K_G^*$  to be the canonical homomorphism given by:  $\bar{\mu}([x/y]) = \mu(x)(\mu(y))^{-1} = \mu(x)$ . Given  $(X, A)$  in  $(C_1)$ , define  $\bar{\mu}: U_G^*(X, A)[\wedge^{-1}] = U_G^*(X, A) \otimes U_G^*[\wedge^{-1}] \rightarrow K_G^*(X, A)$  to be  $\mu \otimes \bar{\mu}: U_G^*(X, A) \otimes U_G^*[\wedge^{-1}] \rightarrow K_G^*(X, A)$ . Q.E.D.

Definition 2.3.4: Let  $V$  be a  $G$ -module .

Define  $\tilde{\rho} \in U_G^2(P(V)) [\wedge^{-1}]$  to be the image of the natural element  $\rho \in U_G^2(P(V))$  ( § 2.2) under the homomorphism:

$$U_G^*(P(V)) \longrightarrow U_G^*(P(V)) \otimes_{U_G^*} U_G^* [\wedge^{-1}] \text{ sending : } \\ x \longrightarrow x \otimes 1 .$$

Given a  $G$ -line bundle  $L$  over a compact  $G$ -space  $X$ , define  $\tilde{\rho}_L$

to be the image of  $\rho_L \in U_G^2(X)$  under the homomorphism :

$$U_G^*(X) \longrightarrow U_G^*(X) \otimes_{U_G^*} U_G^* [\wedge^{-1}]$$

given by :  $x \longrightarrow x \otimes 1$  .

Lemma 2.3.5:

Given a  $G$ -module  $V$ ,  $\bar{\mu}(\tilde{\rho}) = 1 - H$  in  $K_G(P(V))$  where  $H$  is the Hopf bundle over  $P(V)$  and  $\bar{\mu}: U_G^*(P(V)) [\wedge^{-1}] \longrightarrow K_G^*(P(V))$  is the natural homomorphism.

Proof:

By definition of  $\bar{\mu}$ ,  $\bar{\mu}(\tilde{\rho}) = \bar{\mu}(\rho)$  .  $\rho$  is represented by :  $i: P(V)^+ \longrightarrow M_1(G) = P(V \oplus 1)$  where  $i|P(V)$  is inclusion: Hence  $\mu(\rho)$  is the image of  $\bar{\lambda}^1$  in  $\tilde{O}(M_1(G))$  (see § 2.1) under the composition:

$$\tilde{O}(M_1(G)) \xrightarrow{(1)} K_G(P(V \oplus 1)) \xrightarrow{i^*} K_G(P(V)^+) = K_G(P(V))$$

where (1) is the natural map and  $i: P(V) \longrightarrow P(V \oplus 1)$  is inclusion.

By Cor 2.1.6, (1) sends  $\bar{\lambda}^1$  to  $1 - H_1$  where  $H_1$  is the Hopf bundle over  $P(V \oplus 1)$ . Hence  $\mu(\rho) = i^*(1 - H_1) = 1 - H$  in  $K_G(P(V))$  .

Q.E.D.

Cor 2.3.6:

Suppose  $L$  is a  $G$ -line bundle over a compact  $G$ -space  $X$ . Then

$$\bar{\mu}(\bar{\rho}_L) = 1 - L \text{ in } K_G^*(X),$$

Proof:

By the naturality of  $\bar{\mu}$  and  $\bar{\rho}_L$ , and Lemma 2.3.5. Q.E.D.

§ 2.4 :  $U_G^*(-) [\wedge^{-1}]$  - theory Of Projective Spaces:

We assume the following:

Th 2.4.1:

Let  $L_1, \dots, L_n$  be  $G$ -line bundles over a compact  $G$ -space  $X$  and let  $H$  be the Hopf bundle over  $P(L_1 \oplus L_2 \oplus \dots \oplus L_n)$ .

Then  $K_G^*(P(L_1 \oplus \dots \oplus L_n))$  is a free  $K_G^*(X)$ -module generated by  $1, 1-H, (1-H)^2, \dots, (1-H)^{n-1}$ .  $H$  satisfies the relation

$(H - L_1)(H - L_2) \dots (H - L_n) = 0$  and the image of the Thom class  $\gamma$  in  $K_G^*(P_n, P_{n-1})$  under the natural homomorphism

$$K_G^*(P_n, P_{n-1}) \xrightarrow{j!} K_G^*(P_n) \text{ is } (H - L_1)(H - L_2) \dots (H - L_{n-1}),$$

where  $P_m = P(L_1 \oplus L_2 \oplus \dots \oplus L_m)$  ( $1 \leq m \leq n$ ) ( $P_n/P_{n-1}$  can be

identified with the Thom space of  $L_1 \oplus L_2 \oplus \dots \oplus L_{n-1}$ ).

For the proof, see Segal [20] Proposition 3.9 or [4].

Proposition 2.4.2:

Let  $G$  be abelian, and  $V$  a  $G$ -module. Decompose  $V$  in the form

$V = L_1 \oplus \dots \oplus L_n$  ( $\dim L_1 = 1$ ) Let  $H$  be the Hopf bundle over

$P(V)$ .  $U_G^*(P(V)) [\wedge^{-1}]$  is freely generated over  $U_G^* [\wedge^{-1}]$  by

$$1, \bar{\rho}_H, \bar{\rho}_H^2, \dots, \bar{\rho}_H^{n-1}.$$

Proof:

By induction. It is obviously true for  $n = 1$ . Denote

$P(L_1 \oplus \dots \oplus L_m)$  by  $P_m$  ( $1 \leq m \leq n$ ), the Hopf bundle over  $P_m$  by  $H_m$ ,

and  $\bar{\rho}_{H_m}$  by  $\bar{\rho}_m$ . Suppose  $U_G^*(P_m)[\wedge^{-1}]$  is freely generated

over  $U_G^*[\wedge^{-1}]$  by  $1, \bar{\rho}_m, \dots, \bar{\rho}_m^{m-1}$ . Let  $i: P_m \hookrightarrow P_{m+1}$

be inclusion.  $i^*(\bar{\rho}_{m+1}) = \bar{\rho}_m$  and so the cohomology exact

sequence gives us a short exact sequence (s.e.s.):

$$0 \rightarrow U_G^*(P_{m+1}, P_m)[\wedge^{-1}] \xrightarrow{j^*} U_G^*(P_{m+1})[\wedge^{-1}] \xrightarrow{i^*} U_G^*(P_m)[\wedge^{-1}] \rightarrow 0.$$

Since  $U_G^*(P_m)[\wedge^{-1}]$  is free over  $U_G^*[\wedge^{-1}]$ , it is also split.

By the Thom isomorphism theorem (Th 1.4.5),  $U_G^*(P_{m+1}, P_m)[\wedge^{-1}]$

is a free  $U_G^*[\wedge^{-1}]$ -module generated by the Thom class

$t \neq t(L_1 \oplus \dots \oplus L_m)$  (by this we mean the element  $t_m \otimes 1$  in

$U_G^*(P_{m+1}, P_m) \otimes_{U_G^*} U_G^*[\wedge^{-1}]$  where  $t_m = t_{L_1 \oplus \dots \oplus L_m}$  (see § 1.4)).

Hence  $U_G^*(P_{m+1})[\wedge^{-1}]$  is freely generated by:

$1, \bar{\rho}_{m+1}, \bar{\rho}_{m+1}^2, \dots, \bar{\rho}_{m+1}^{m-1}, j^*(t)$ . In particular,  $\exists$

elements  $a_0, a_1, \dots, a_{m-1}, b \in U_G^*[\wedge^{-1}]$  such that:

$$\prod_{k=1}^m (\bar{\rho}_{m+1} - \bar{\rho}_{L_k}) = \sum a_s \bar{\rho}_{m+1}^s + b \cdot j^*(t) \dots \quad (i)$$

If we take  $\bar{\mu}$  of each side of (i), we get

$$(-)^m \prod_{k=1}^m (H_{m+1} - L_k) = \sum (-)^s \bar{\mu}(a_s) (H_{m+1} - 1)^s + \bar{\mu}(b) \bar{\mu} j^*(t),$$

(Cor 2.3.5). By considering the commutative diagram:

$$\begin{array}{ccc} U_G^* (P_{m+1}, P_m) [\wedge^{-1}] & \xrightarrow{j^*} & U_G^* (P_{m+1}) [\wedge^{-1}] \\ \downarrow \bar{\mu}' & & \downarrow \bar{\mu} \\ K_G^* (P_{m+1}, P_m) & \xrightarrow{j!} & K_G^* (P_{m+1}) \end{array}$$

and Th 2.4.1, we deduce :  $\bar{\mu}_{j^*} (t) = j! \bar{\mu}'(t)$

i.e.  $\bar{\mu}_{j^*} (t) = j! (\tau) \quad (\text{Lemma 2.1.7})$

$$= \prod_{k=1}^m (H_{m+1} - L_k) \quad (\text{Th 2.4.1}). \quad \text{Thus}$$

$$(-)^m \prod_{k=1}^m (H_{m+1} - L_k) = \sum (-)^s \bar{\mu}(a_s) (H_{m+1} - 1)^s + \bar{\mu}(b) \prod_{k=1}^m (H_{m+1} - L_k).$$

Since  $1, 1-H_{m+1}, (1-H_{m+1})^2, \dots, (1-H_{m+1})^{m-1}$  &  $\prod_{k=1}^m (H_{m+1} - L_k)$

generate  $K_G^* (P_{m+1})$  freely over  $RG$  (Th 2.4.1), it follows that

$\bar{\mu}(b) = 1$  or  $-1$ . Hence  $b$  is an invertible element of

$U_G^* [\wedge^{-1}]$  and  $U_G^* (P_{m+1}) [\wedge^{-1}]$  is a free  $U_G^* [\wedge^{-1}]$ -module

generated by :  $1, \bar{e}_{m+1}, \bar{e}_{m+1}^2, \dots, \bar{e}_{m+1}^{m-1}, \& \prod_{k=1}^m (\bar{e}_{m+1} - \bar{e}_{L_k})$

i.e. we may also take as generators :

$$1, \bar{e}_{m+1}, \bar{e}_{m+1}, \dots, \bar{e}_{m+1}^m. \quad \text{Q.E.D.}$$

### Cor 2.4.3:

Let  $E = \mathbb{X} \times V$  be a trivial  $G$ -vector bundle over the compact  $G$ -space

$X$  ( $G$  abelian). Let  $H_0$  be the Hopf bundle over  $P(E)$ .

Then  $U_G^* (P(E)) [\wedge^{-1}]$  is a free  $U_G^* (X) [\wedge^{-1}]$ -module generated by

$$1, \bar{e}_{H_0}, \bar{e}_{H_0}^2, \dots, \bar{e}_{H_0}^{1V1-1}.$$

Proof:-

Again by induction. Put  $V = L_1 \oplus \dots \oplus L_n$  ( $\dim L_1 = 1$ ). Let  $\pi_m : X \times P_m \rightarrow P_m$  be projection onto the 2nd factor. Suppose the result is true for  $X \times P_m$ . We have a commutative diagram:

$$\begin{array}{ccc} U_G^*(X \times P_{m+1}, X \times P_m) [\wedge^{-1}] & \xrightarrow{j^*} & U_G^*(X \times P_{m+1}) [\wedge^{-1}] \\ \uparrow \pi^* & & \uparrow \pi_{m+1}^* \\ U_G^*(P_{m+1}, P_m) [\wedge^{-1}] & \xrightarrow{j_2^*} & U_G^*(P_{m+1}) [\wedge^{-1}] \end{array}$$

With the notation used in the proof of the previous proposition, we have shown that:

$$\prod_{k=1}^m (\bar{\rho}_{m+1} - \bar{\rho}_{L_k}) = \sum a_s \bar{\rho}_{m+1}^s + b \cdot j_2^*(t)$$

where  $b$  is invertible in  $U_G^*[\wedge^{-1}]$ . Now  $\pi_{m+1}^*(\bar{\rho}_{m+1}) = \bar{\rho}_{m+1} =$  the natural element in  $U_G^2(X \times P_{m+1})[\wedge^{-1}]$ . On the other hand,  $\pi^*(t) = \bar{t}$  = the Thom class in  $U_G^*(X \times P_{m+1}, X \times P_m) [\wedge^{-1}]$ .

$$\text{Hence } \prod_{k=1}^m (\bar{\rho}_{m+1} - \rho_{L_k}) = \sum a_s \bar{\rho}_{m+1}^s + b j^*(\bar{t}).$$

By the same argument as in the proof of Proposition 2.4.2, we conclude that  $U_G^*(X \times P_{m+1})[\wedge^{-1}]$  is freely generated by:

$$1, \bar{\rho}_{m+1}, \bar{\rho}_{m+1}^2, \dots, \bar{\rho}_{m+1}^m. \quad \text{Q.E.D.}$$

Proposition 2.4.4:

Let  $G$  be abelian. Suppose  $E$  is an  $n$ -dimensional  $G$ -vector bundle over the compact  $G$ -space  $X$ . Then  $U_G^*(P(E))[\wedge^{-1}]$  is a free

$U_G^*(X)$  - module generated by :  $1, \bar{\rho}, \dots, \bar{\rho}^{n-1}$  where  
 $\rho \in U_G^2(P(E))[\wedge^{-1}]$  is the natural element .

Proof:-

Given  $x \in X$ , let  $G_x$  be the stabilizer. The fibre  $E_x$  is a  $G_x$ -module. Because  $G$  is abelian, one can regard it as the restriction of a  $G$ -module. Then  $E|_{Gx} = Gx \times E = (G/G_x) \times E_x$ . So  $E$  and  $X \times E$  are isomorphic on the orbit  $Gx$  and hence on a closed neighbourhood  $F$  of it. Let  $F_0$  be a  $G$ -neighbourhood of  $Gx$  and a closed subspace of  $F$ . By Cor 2.4.3,  $U_G^*(P(F_0 \times E_x))[\wedge^{-1}]$  is a free  $U_G^*(F_0)[\wedge^{-1}]$  - module generated by  $1, \bar{\rho}, \dots, \bar{\rho}^{n-1}$  where  $\bar{\rho} \in U_G^2(P(F_0 \times E_x))[\wedge^{-1}]$  is the natural element. The isomorphism  $E|_{F_0} \cong F_0 \times E_x$  induces an isomorphism of the projective bundles  $\beta: P(E|_{F_0}) \cong P(F_0 \times E_x)$ , and of the Hopf bundles over them. Therefore, the natural elements  $\bar{\rho} \in U_G^2(P(E|_{F_0}))[\wedge^{-1}]$  and  $\bar{\rho} \in U_G^2(P(F_0 \times E_x))[\wedge^{-1}]$  correspond to one another under  $\beta$  i.e.  $U_G^*(P(E|_{F_0}))[\wedge^{-1}]$  is a free  $U_G^*(F_0)[\wedge^{-1}]$  - module generated by  $1, \bar{\rho}, \dots, \bar{\rho}^{n-1}$  .

By Proposition 2.3.1 and Proposition 1.6.3, the result now follows:

Q.E.D.

Chapter 3 : Applications

3.1 : Characteristic Classes

Let  $h_G^* = \{h_G^n\}$  be a multiplicative equivariant cohomology theory on  $(c_1)$ . Suppose that  $\forall$   $G$ -module  $V$ ,  $\exists$  a natural element  $\sigma \in h_G^2(P(V))$  such that if  $i: P(V_1) \rightarrow P(V_2)$  is inclusion, then  $i^*(\sigma_2) \overset{\text{equals}}{=} \sigma_1$  where  $\sigma_i \overset{\text{equals}}{=}$  the natural element  $\sigma \in h_G^2(P(V_i))$ . Assign to every  $G$ -line bundle  $L$  over a compact  $G$ -space  $X$ , an element  $c_1(L) \in h_G^2(X)$  by the requirements : (a) if  $H$  is the Hopf bundle over  $P(V)$ ,  $c_1(H) = \sigma \in h_G^2(P(V))$  and (b) if  $f: X \rightarrow P(W)$  is a classifying map for  $L$ , and  $f^*: h_G^2(P(W)) \rightarrow h_G^2(X)$  the induced homomorphism, then  $c_1(L) = f^*(c_1(H))$  where  $H$  is the Hopf bundle over  $P(W)$ . That  $c_1(L)$  does not depend on the choice of  $f$  follows as in Definition 2.2.1. If  $L' = f^*(L)$ ,  $c_1(L') = f^*(c_1(L))$ .

Proposition 3.1.1:

Let  $G$  be abelian. Suppose that  $\forall$  compact  $G$ -space  $X$  and  $\forall$   $G$ -module  $V$ ,  $h_G^*(P(V \times X))$  is a free  $h_G^*(X)$  - module generated by  $1, \sigma, \sigma^2, \dots, \sigma^{1V1-1}$  where  $\sigma$  = the natural element in  $h_G^2(P(V \times X))$ . Then we can assign to every  $n$ -dimensional  $G$ -vector bundle  $E$  over a compact  $G$ -space  $X$  a Chern class:  

$$c(E) = 1 + c_1(E) + c_2(E) + \dots + c_n(E) \quad , \quad c_i(E) \in h_G^{2i}(X).$$
 Moreover, if  $E' = f^*(E)$ ,  $c(E') = f^*(c(E))$ .



Proof:

If  $1E1 = 1$ , define  $c_1(E)$  as above.

Suppose  $n = 1E1 > 1$ . Let  $\sigma = c_1(H)$  where  $H$  is the Hopf bundle over  $P(E)$ . Because  $G$  is abelian,  $E$  is locally decomposable ([20] Proposition 3.7). By Proposition 1.6.3, it follows as in the proof of Proposition 2.4.4, that  $h_G^*(P(E))$  is a free  $h_G^*(X)$  - module generated by :

$1, \sigma, \sigma^2, \dots, \sigma^{n-1}$ . Define  $c_1(E), \dots, c_n(E)$  to be the unique elements in  $h_G^*(X)$  satisfying :

$$(3.1.2) \quad \sum_{i=0}^n (-1)^i c_i(E) \sigma^{n-i} = 0. \quad \text{Q.E.D.}$$

Suppose that  $h_G^*$  satisfies the additional property :-

(\*) given based  $G$ -maps of compact based  $G$ -spaces  $f: X \rightarrow X^1$ ,  $g: Y \rightarrow Y^1$  and  $a \in \tilde{h}_G^i(X^1)$ ,  $b \in \tilde{h}_G^j(Y^1)$ , then  $(f \wedge g)^*(a \otimes b) = f^* a \otimes g^* b$ .

In that case,

Proposition 3.1.3:

The Chern classes defined by (3.1.2) satisfy the Whitney Product Theorem i.e. given an  $m$ -dimensional  $G$ -vector bundle  $E_1$  and an  $n$ -dimensional  $G$ -vector bundle  $E_2$  over the compact  $G$ -space  $X$ , then  $c(E_1 \oplus E_2) = c(E_1) \cdot c(E_2)$ .

Proof:

Put an invariant metric on  $E_1 \oplus E_2$ . Then  $P(E_1 \oplus E_2) = S(E_1 \oplus E_2)/U(1)$

where  $S(E_1 \oplus E_2) = \{(e_1, e_2) \mid \|e_1\|^2 + \|e_2\|^2 = 1\}$ .

Let A & B be the closed subspaces of  $P(E_1 \oplus E_2)$  defined by :

$$A = \{[e_1, e_2] : \|e_2\|^2 \leq \frac{1}{2}\}, B = \{[e_1, e_2] : \|e_1\|^2 \leq \frac{1}{2}\}.$$

Then A properly deformation retracts onto  $P(E_1)$ , B properly deformation retracts onto  $P(E_2)$ , and  $P(E_1 \oplus E_2) = A \cup B$ . Let

$\sigma$  be the natural element in  $h_G^2(P(E_1 \oplus E_2))$ . Then

$$\chi = \sum_{i=0}^m (-)^i c_i(E_1) \sigma^{m-i} \text{ restricts to } 0 \text{ in } h_G^*(A). \text{ Let}$$

$j_1 : (A \cup B, \emptyset) \rightarrow (A \cup B, A)$  be the inclusion.  $\exists \chi_1 \in h_G^*(A \cup B, A)$  such that  $j_1^*(\chi_1) = \chi$ . Similarly, if  $y = \sum_{k=0}^n (-)^k c_k(E_2) \sigma^{n-k}$

and  $j_2 : (A \cup B, \emptyset) \rightarrow (A \cup B, B)$  is the inclusion,  $\exists y_1 \in h_G^*(A \cup B, B)$  such that  $j_2^*(y_1) = y$ . The element  $\chi_1 \otimes y_1$  which

$\in h_G^*((A \cup B) \times (A \cup B), A \times (A \cup B) \cup (A \cup B) \times B)$  then has

$$\beta^*(\chi_1 \otimes y_1) = \chi \otimes y \text{ where}$$

$$\beta : (A \cup B) \times (A \cup B) \hookrightarrow ((A \cup B) \times (A \cup B), A \times (A \cup B) \cup (A \cup B) \times B)$$

(because multiplication satisfies the property (\*)) .

Therefore,  $\chi \cdot y = 0$  in  $h_G^*(A \cup B)$  since  $h_G^*(Y, Y) = 0$  for any  $(Y, Y)$  i.e.

$$\left( \sum_{i=0}^m (-)^i c_i(E_1) \sigma^{m-i} \right) \left( \sum_{k=0}^n (-)^k c_k(E_2) \sigma^{n-k} \right)$$

$$= 0 \text{ in } h_G^*(E_1 \oplus E_2). \text{ Also}$$

$$\sum_{i=0}^{m+n} (-)^i c_i(E_1 \oplus E_2) \sigma^{m+n-i} = 0. \text{ Hence by}$$

$$\text{Proposition 3.1.1., } c_k(E_1 \oplus E_2) = \sum_{i+j=k} c_i(E_1) \cdot c_j(E_2). \quad \text{Q.E.D.}$$

Remark:

If  $h_G^*$  admits Chern classes satisfying :  $c_1(E_1 \oplus E_2) = c_1(E_1) + c_1(E_2)$ , then for all compact connected G-spaces X, we can define an additive homomorphism:

$$c_1 : K_G(X) \longrightarrow h_G^2(X)$$

by :  $[E] \longrightarrow c_1(E)$ .

As examples of equivariant cohomology theories admitting Chern classes:

Cor 3.1.4:

Let G be abelian. Given an n-dimensional G-vector bundle E over a compact G-space X, let  $\sigma = \bar{\rho}_H$  (Def.2.3.4) where H is the Hopf bundle over P(E). We can assign to E a class  $cf(E) = 1 + cf_1(E) + \dots + cf_n(E)$ ,  $cf_i(E) \in U_G^{2i}(X)[\wedge^{-1}]$ , given by:

$$\sum_{i=0}^n (-1)^i cf_i(E) \bar{\rho}_H^{n-i} = 0 \quad (*)$$

cf is natural w.r.t. bundle maps and  $cf(E_1 \oplus E_2) = cf(E_1) \cdot cf(E_2)$

Proof:

From propositions 2.4.4, 3.1.1, 3.1.3.

Q.E.D.

Cor 3.1.5:

Let E & X be as above. We can assign to E a class

$\overline{cf}(E) = 1 + \overline{cf}_1(E_1) + \dots + \overline{cf}_n(E)$ ,  $\overline{cf}_i \in K_G^{2i}(X)$ , defined by

$$\sum_{i=0}^n (-1)^i \overline{cf}_i(E) (1-H)^{n-i} = 0 \quad (**)$$

Proposition 3.1.6:

Let  $\bar{\mu} : U_G^*(X)[\wedge^{-1}] \longrightarrow K_G^*(X)$  be the canonical map. Then

$$\overline{cf}(E) = \bar{\mu}(cf(E))$$

Proof:

Let  $\bar{\mu}': U_G^*(P(E))[\wedge^{-1}] \rightarrow K_G^*(P(E))$ . By Cor 2.3.6,

$\bar{\mu}'(\bar{\rho}_H) = 1 - H$ . Taking  $\bar{\mu}'$  of (\*) and using the fact that  $\bar{\mu}$  is natural and multiplicative, we get:

$$\sum_{i=0}^n (-1)^i \bar{\mu}(cf_i(E)) (1-H)^{n-i} = 0. \text{ Comparing}$$

with (\*\*),  $\bar{cf}_i(E) = \bar{\mu}(cf_i(E))$ . Q.E.D.

§ 3.2 : An embedding of  $K_G(X, A)$  in  $U_G^0(X, A)[\wedge^{-1}]$ :

From the results of § 3.1, we can define,  $\forall$  compact connected G-space X, a homomorphism :

$$c_1 : K_G(X) \longrightarrow U_G^2(X)[\wedge^{-1}]$$

by:  $c_1([E] - [E^1]) = cf_1(E) - cf_1(E^1)$ . We assume that G is abelian.

Lemma 3.2.1:

Let X be a compact connected G-space with base point (G abelian).

The composition:

$$\begin{aligned} \tilde{K}_G(X) &\xrightarrow{c_1} \tilde{U}_G^2(X)[\wedge^{-1}] \xrightarrow{\bar{\mu}} \tilde{K}_G(X) \\ \text{is} = -\text{id} : \tilde{K}_G(X) &\longrightarrow \tilde{K}_G(X). \end{aligned}$$

Proof:

By induction on the dimension of E over X. If L is a G-line bundle, then  $\bar{\mu}(cf_1(L)) = \bar{cf}_1(L) = 1 - L$  (Proposition 3.1.6) i.e.  $\bar{\mu}(cf_1(L - 1)) = -(L - 1)$ .

Suppose that  $\forall$   $k$ -dimensional  $G$ -vector bundle  $E$  over  $X$ ,

$\overline{cf}_1(E - k) = - (E - k) \ (\overline{cf} = \mu \circ cf)$ . Let  $E_1$  have dimension  $k+1$ . Let  $p: P(E_1) \rightarrow X$  be the projection, and  $H$  the Hopf bundle over  $P(E_1)$ .  $p^*(E_1) = H \oplus E$  and by ~~Th 2.4.1~~  $p^*: K_G(X) \rightarrow K_G(P(E_1))$  is injective. Hence  $p^* \overline{cf}_1(E_1 - \overline{k+1}) = \overline{cf}_1(H + E - \overline{k+1}) = - (H + E - \overline{k+1})$  (by the induction hypothesis)  $= - p^*(E_1 - \overline{k+1})$  i.e.  $\overline{cf}_1(E_1 - \overline{k+1}) = - (E - \overline{k+1})$ .

The induction is complete.

Q.E.D.

Given a compact  $G$ -space  $X$  with base point, form  $S^2 \wedge X$ . It is a connected space. So we can define a homomorphism:

$$\underline{c}_0: \widetilde{K}_G(X) \longrightarrow \widetilde{U}_G^0(X) [\wedge^{-1}] \quad (G \text{ abelian})$$

as the composition:

$$\begin{aligned} \widetilde{K}_G(X) &\cong \widetilde{K}_G^2(S^2 \wedge X) \cong \widetilde{K}_G(S^2 \wedge X) \xrightarrow{\underline{c}_1} U_G^2(S^2 \wedge X) [\wedge^{-1}] = \\ &\cong \widetilde{U}_G^0(X) [\wedge^{-1}] \end{aligned}$$

where all the isomorphisms are the canonical ones.

### Cor 3.2.2:

Let  $G$  be abelian. Given a locally compact  $G$ -space  $X$  and a closed  $G$ -subspace  $A$  of  $X$ ,  $\exists$  an additive homomorphism

$$\underline{c}_0: K_G(X, A) \longrightarrow U_G^0(X, A) [\wedge^{-1}]$$

such that  $\bar{\mu} \circ \underline{c}_0 = - \text{id}: K_G(X, A) \longrightarrow K_G(X, A)$ .

Proof:

Given a compact G-space X with base point, the diagram:

$$\begin{array}{ccc}
 \tilde{K}_G(X) & \xleftarrow{\bar{\mu}} & \tilde{U}_G^0(X)[\wedge^{-1}] \\
 \downarrow \cong & & \uparrow \cong \\
 \tilde{K}_G^2(S^2 \wedge X) & \xleftarrow{\bar{\mu}} & \tilde{U}_G^2(S^2 X)[\wedge^{-1}] \\
 \parallel & & \nearrow \subseteq \\
 \tilde{K}_G(S^2 \wedge X) & & 
 \end{array}$$

is commutative. Now apply Lemma 3.2.1.

Q.E.D.

### § 3.3 : $U_G^*(-)[\wedge^{-1}]$ of Grassmannians :

#### Proposition 3.3.1 :

Let  $E$  be  $\wedge$   $G$ -vector bundle over a compact  $G$ -space  $X$  and let  $F(E)$  be the flag bundle of  $E$ . Let  $L_1, \dots, L_n$  be the natural line bundles over  $F(E)$ . Then  $K_G^*(F(E))$  is a free  $K_G^*(X)$ -module generated by a finite number of elements of the form  $L_1^{k_1} L_2^{k_2} \dots L_n^{k_n}$  where  $k_i \in \mathbb{Z}^+ \forall i$ .

#### Proof:

If  $n=1$ ,  $F(E) = X$  and the proposition is true. Suppose the proposition is true for  $n-1$  ( $n > 1$ ) and let  $E$  have dimension  $n$ . Let  $P(E)$  be the projective bundle of  $E$ ,  $H$  the Hopf bundle over  $P(E)$  and  $p: P(E) \rightarrow X$  the natural projective. Define  $E'$  over  $P(E)$  by the relation :

$$p^*(E) = H \oplus E'$$

There is a natural identification :  $F(E) = F(E')$

by sending the flag  $\theta = M_0, M_1, M_2, \dots, M_n = E_x$  ( $x \in X$ )

of linear subspaces of  $E_x$  to the flag  $M_0, M_2/M_1, M_3/M_1, \dots,$

$M_n/M_1 = E'_{M_1}$ . Under this identification, the natural line

bundles  $L_2, \dots, L_n$  over  $F(E)$  correspond to the natural line bundles  $L'_1, L'_2, \dots, L'_{n-1}$  over  $F(E')$ , and  $L_1$  over

$F(E)$  corresponds to  $H$  lifted to  $F(E')$ . Since  $K_G^*(P(E))$  is

freely generated over  $K_G^*(X)$  by  $1, H, \dots, H^{n-1}$  and the diagram :

$$\begin{array}{ccc}
 F(E) & = & F(E') \\
 \downarrow & & \downarrow \\
 X & \longleftarrow & P(E)
 \end{array}$$

is commutative, the proposition is then true for  $\dim E = n$ .

By induction, it is true  $\forall n$ .

Q.E.D.

Cor 3.3.2:

Let  $V$  be a  $G$ -module,  $L$  a 1 - dim<sup>L</sup>  $G$ -module. The natural map:

$$K_G^*(F(V \oplus L)) \xrightarrow{i!} K_G^*(F(V))$$

is an epimorphism.

Proof:

Follows from the above proposition because each of the natural line bundles over  $F(V)$  is the restriction of a natural one over  $F(V \oplus L)$ .

Q.E.D.

Lemma 3.3.3:

Let  $V$  be a  $G$ -module,  $G_k(V)$  the Grassmann manifold (of  $k$ -dim. subspaces) of  $V$ . Let  $F(V)$  be the flag manifold of  $V$ .  $\exists$  an embedding  $K_G^*(G_k(V)) \xrightarrow{y} K_G^*(F(V))$  with a natural left inverse:

$$Y_! : K_G^*(F(V)) \longrightarrow K_G^*(G_k(V))$$

Proof:

Let  $p(V)$  be the associated principal bundle. Because

$$U(k) \times U(n-k) \text{ acts freely on } p(V) \quad (n=1V1), \text{ then } K_G^*(G_k(V)) = K_G^*(p(V)/U(k) \times U(n-k)) = K_G^* \times U(k) \times U(n-k) (p(V))$$



([20]). Similarly if  $T$  is a maximal torus in  $U(n)$ , then  $K_G^*(F(V)) = K_{G \times T}^*(p(V))$ . It is shown by Atiyah ([2]) that the restriction homomorphism :

$$K_{G \times U(k) \times U(n-k)}^*(p(V)) \longrightarrow K_{G \times T}^*(p(V))$$

has a functional left inverse. Hence the result.

Q.E.D.

Lemma 3.3.4:

Let  $V$  be a  $G$ -module,  $L$  a 1-dim.  $G$ -module. The  $G$ -space  $G_k(V \oplus L)/G_k(V)$  is  $G$ -homeomorphic to the Thom space of a  $G$ -vector bundle (of a certain dimension) over  $G_{k-1}(V)$ . This is proved in the ordinary case,  $G=e$ , by S. Hoggar [13]. His proof is still valid in this general setting.

Proposition 3.3.5:

- (i) If  $L$  is a 1 - dim  $G$ -module, and  $V$  is a  $G$ -module, the natural homomorphism :-

$$K_G^*(G_k(V \oplus L)) \xrightarrow{j^!} K_G^*(G_k(V))$$

is epi.

- (ii) Given a decomposable  $G$ -module  $V$ , then  $K_G^*(G_k(V))$  is a finitely-generated free  $RG$  - module.

Proof:

- (i) By Cor 3.3.2, the natural homomorphism

$$K_G^*(F(V \oplus L)) \xrightarrow{i^!} K_G^*(F(V)). \text{ is epi. The diagram:}$$

$$\begin{array}{ccc}
 K_G^*(G_k(V \oplus L)) & \xrightarrow{j^!} & K_G^*(G_k(V)) \\
 \Upsilon_! \uparrow & & \uparrow \Upsilon_! \\
 K_G^*(F(V \oplus L)) & \xrightarrow{i^!} & K_G^*(F(V))
 \end{array}$$

is commutative. Thus  $j^! \circ \Upsilon_!$  is an epimorphism. This implies  $j^!$  is epi.

(ii) Follows from (i) by using the Thom isomorphism theorem for  $K_G$  - theory ([2]), and Lemma 3.3.4. Q.E.D.

Proposition 3.3.6:

Let  $V$  be a  $G$ -module ( $G$  abelian). Then  $U_G^*(G_k(V))[\wedge^{-1}]$  is freely generated over  $U_G^*[\wedge^{-1}]$  by a finite number of elements  $x_1, \dots, x_t$  such that  $\tilde{\mu}(x_1), \tilde{\mu}(x_2), \dots, \tilde{\mu}(x_t)$  generate  $K_G^*(G_k(V))$  freely over  $RG$ .

Proof:

By induction on  $k$ . If  $k = 1$ ,  $G_1(V) = P(V)$  and the result follows from Proposition 2.

Suppose it is true for  $k - 1$ , and we want to prove it for  $k(k \geq 2)$ . We proceed by induction on  $|V|$ . For  $|V| = k$ ,  $G_k(V) = \text{pt.}$  and the result is true. Suppose the proposition has been proved for  $|V| = n$ . Let  $L$  be a 1 - dim.  $G$ -module.

We have a commutative diagram:

$$\begin{array}{ccccc}
 U_G^*(G_k(V \oplus L), G_k(V))[\wedge^{-1}] & \rightarrow & U_G^*(G_k(V \oplus L))[\wedge^{-1}] & \xrightarrow{i^*} & U_G^*(G_k(V))[\wedge^{-1}] \\
 \tilde{\mu} \downarrow & & \tilde{\mu} \downarrow & & \tilde{\mu} \downarrow \\
 K_G^*(G_k(V \oplus L), G_k(V)) & \longrightarrow & K_G^*(G_k(V \oplus L)) & \xrightarrow{i^!} & K_G^*(G_k(V))
 \end{array}$$

By hypothesis,  $U_G^*(G_k(V))$  is freely generated by  $x_1, \dots, x_l$  such that  $y_1, \dots, y_l$  ( $y_i = \bar{\mu}(x_i)$ ) generate  $K_G^*(G_k(V))$  freely over  $RG$ . By Proposition 3.3.5 (i),  $y_i$  can be pulled back to an element  $y_i' \in K_G^*(G_k(V \oplus L))$ . Since

$\bar{\mu}: U_G^*(G_k(V \oplus L))[\wedge^{-1}] \longrightarrow K_G^*(G_k(V \oplus L))$  is epi (Cor 3.2.2),

$y_i'$  can be pulled back to some element in  $U_G^*(G_k(V \oplus L))[\wedge^{-1}]$ .

In particular,  $y_1' = \bar{\mu}(t)$ , some  $t$ . Let  $z_1 = i^*(t)$ . Then

$\bar{\mu}(z_1) = \bar{\mu}i^*(t) = i^! \bar{\mu}(t) = y_1$ . By the induction

hypothesis,  $\exists a_1, \dots, a_l \in U_G^*[\wedge^{-1}]$  such that  $z_1 = \sum a_i x_i$ .

Hence  $\bar{\mu}(z_1) = \sum \bar{\mu}(a_i) y_i$  i.e.  $y_1 = \sum \bar{\mu}(a_i) y_i$ . This

implies  $\bar{\mu}(a_1) = 1$ . But  $a_1 = [s/s]$  where  $\bar{\mu}(s) = 1$ .

Therefore,  $\bar{\mu}(s) = 1$  i.e.  $s$  is invertible in  $U_G^*[\wedge^{-1}]$ ,

and so is  $a_1$ . Hence if we replace  $x_1$  by  $z_1$ , we still have

a basis  $z_1, x_2, x_3, \dots, x_l$  of  $U_G^*(G_k(V))[\wedge^{-1}]$ .

Proceeding in this way, we obtain a basis  $z_1, \dots, z_l$  of

$U_G^*(G_k(V))[\wedge^{-1}]$  such that  $z_j = i^*(t_j) \forall j$ .

Hence  $i^*: U_G^*(G_k(V \oplus L))[\wedge^{-1}] \longrightarrow U_G^*(G_k(V))[\wedge^{-1}]$  is epi.

Since the latter module is free, the sequence:

$$U_G^*(G_k(V \oplus L), G_k(V))[\wedge^{-1}] \rightarrow U_G^*(G_k(V \oplus L))[\wedge^{-1}] \xrightarrow{i^*} U_G^*(G_k(V))[\wedge^{-1}]$$

is a split s.e.s.

Now using the Thom isomorphism theorem (Th 1.4.5), Lemma 3.3.4,

Lemma 2.1.7, and the induction hypothesis, we see that the

result is valid for  $G_k(V \oplus L)$ . But  $G$  is abelian and so every

$G$ -module is decomposable. Hence the result.

Q.E.D.

### § 3.4 : Equivariant K-theory and Cobordism (G abelian):

We are now in a position to relate  $K_G$  - theory to  $U_G^*$  - theory for G abelian. The non-abelian case is also true but needs more work. The next chapter will be concerned mainly with that.

Regard  $RG$  as a  $U_G^* (\Lambda^{-1})$  - module via the homomorphism:

$$\bar{\mu} : U_G^* (\Lambda^{-1}) \longrightarrow RG \quad (K_G^1(\text{pt}) = 0). \quad \text{Given } (X, A) \text{ in } (C_1),$$

define :

$$L_G^0(X, A) = U_G^{\text{ev}}(X, A) [\Lambda^{-1}] \otimes_{U_G^* [\Lambda^{-1}]}^{RG} , \quad L_G^1(X, A) = U_G^{\text{od}}(X, A) \otimes_{U_G^* [\Lambda^{-1}]}^{RG}$$

$$\text{and } L_G^*(X, A) = L_G^{\text{od}}(X, A) \oplus L_G^1(X, A) \quad \dots \quad (3.4.1)$$

There are natural homomorphisms (i)  $\beta : U_G^*(X, A) [\Lambda^{-1}] \longrightarrow L_G^*(X, A)$

defined by  $\beta(x) = x \otimes 1$ , and (ii)

$$\hat{\mu} : U_G^*(X, A) [\Lambda^{-1}] \otimes_{U_G^* [\Lambda^{-1}]}^{RG} \longrightarrow K_G^*(X, A)$$

with  $\hat{\mu}(x \otimes y) = y \cdot \bar{\mu}(x)$ . Hence there is a commutative diagram:

$$\begin{array}{ccc} U_G^*(X, A) [\Lambda^{-1}] & \xrightarrow{\beta} & L_G^*(X, A) \\ & \searrow \hat{\mu} & \swarrow \hat{\mu} \\ & K_G^*(X, A) & \end{array}$$

Define  $\hat{c}_0 : K_G^*(X, A) \longrightarrow L_G^*(X, A)$  by the composition:

$$K_G^*(X, A) \xrightarrow{c_0} U_G^*(X, A) [\Lambda^{-1}] \xrightarrow{\beta} L_G^*(X, A)$$

Then  $\hat{\mu} \circ \hat{c}_0 = -\text{id} : K_G^*(X, A) \longrightarrow K_G^*(X, A)$ . Hence

$$\hat{\mu} : L_G^*(X, A) \longrightarrow K_G^*(X, A) \text{ is epi.}$$

We can define, too, a  $U_G^*$ -module structure on  $RG$  by means of  $\tilde{\mu}: U_G^* \longrightarrow RG$ . In a similar way,  $U_G^*(X,A) \otimes_{U_G^*} RG$  is  $\mathbb{Z}_2$ -graded via:

$$U_G^*(X,A) \otimes_{U_G^*} RG = U_G^{ev}(X,A) \otimes_{U_G^*} RG \oplus U_G^{od}(X,A) \otimes_{U_G^*} RG.$$

Th 3.4.2 :

Let  $G$  be abelian. Suppose  $X$  is a locally compact  $G$ -space, and  $A$  a closed  $G$ -subspace of  $X$ . The homomorphism:

$$\mu \otimes \text{id}: U_G^*(X,A) \otimes_{U_G^*} RG \longrightarrow K_G^*(X,A)$$

is an isomorphism of  $\mathbb{Z}_2$ -graded rings.

Proof:

$$\begin{aligned} \text{Because of the identification } U_G^*(X,A) \otimes_{U_G^*} RG &= U_G^*(X,A) \otimes_{U_G^*} (U_G^*[\wedge^{-1}] \otimes_{U_G^*} RG) \\ &= (U_G^*(X,A) \otimes_{U_G^*} U_G^*[\wedge^{-1}]) \otimes_{U_G^*[\wedge^{-1}]} RG = U_G^*(X,A)[\wedge^{-1}] \otimes_{U_G^*[\wedge^{-1}]} RG, \end{aligned}$$

it is enough to show that

$$\hat{\mu}: L_G^*(X,A) \longrightarrow K_G^*(X,A)$$

is an isomorphism of  $\mathbb{Z}_2$ -graded rings. We have already shown it is an epimorphism, and so it only remains to prove that it is a monomorphism. Let  $V$  be a  $G$ -module,  $G_k(V)$  the Grassmann manifold of  $k$ -dim. Subspaces of  $V$  and  $M_k(V)$  the Thom space of the standard bundle over  $G_k(V)$ . It follows from Proposition 3.3.6 that  $\hat{\mu}: L_G^*(G_k(V)) \xrightarrow{\cong} K_G^*(G_k(V))$  (See [8] P.60-61).

By the Thom isomorphism theorem for  $U_G^*[\wedge^{-1}]$  - theory, and the commutative diagram (Lemma 2.17):

$$\begin{array}{ccc} L_G^*(G_k(V)) & \xrightarrow{\phi} & L_G^*(M_k(V)) \\ \hat{\mu} \downarrow & & \downarrow \hat{\mu} \\ K_G^*(G_k(V)) & \xrightarrow[\cong]{\phi} & K_G^*(M_k(V)) \end{array}$$

$\hat{\mu}: L_G^*(M_k(V)) \xrightarrow{\cong} K_G^*(M_k(V))$ . This implies the general case by using essentially the argument of Conner-Floyd [8] Theorem (10.1). Q.E.D.

#### Chapter 4 : The Relation of $K_G$ - theory to Cobordism:

We wish to prove Th 3.4.2 for general  $G$ . This is achieved by studying the relation of  $U_{U(n)}^*(-)$  and  $U_T^*(-)$  where  $T$  is a maximal torus of  $U_n$ .

##### § 4.1 : The Relation of $U_G^*(-)$ with Bordism:

Closely related to  $U_G^*$  is the bordism ring  $\mathcal{U}_G^*$  of unitary  $G$ -manifolds (for the definition see T. tom Dieck [11] P.29).

Recall that an element of  $\mathcal{U}_G^{-n}$  ( $n \in \mathbb{Z}^+$ ) is represented by a closed unitary  $G$ -manifold ([11] P.29)  $N^n$  of dimension  $n$ , and

$[N_1^n] + [N_2^n] = [N_1^n \cup N_2^n]$ . Multiplication in  $\mathcal{U}_G^* = \sum \mathcal{U}_G^{-n}$  is induced by cartesian product:  $[N_1^k] \times [N_2^l] = [N_1^k \times N_2^l]$ .

$\mathcal{U}_G^*$  has a unit  $1 \in \mathcal{U}_G^0$  represented by the  $G$ -manifold consisting of a single point.

In the manner of Thom [23], one can define a ring homomorphism (T. tom Dieck [11]).

$$\iota : \mathcal{U}_G^* \longrightarrow U_G^* = U_G^*(pt),$$

as follows. Suppose  $x \in \mathcal{U}_G^{-n}$  is represented by the closed unitary  $G$ -manifold  $M^n$ ,  $n$  even. Embed  $M$  in a  $G$ -module  $V$  (Palais [18]) such that the normal bundle  $N$  of  $M$  in  $V$  receives the correct complex structure as a  $G$ -vector bundle (by the definition of unitary  $G$ -manifold [11]). Now  $N$  can be identified with a  $G$ -tubular neighbourhood of  $M$  in  $V$ . Let  $T, \partial T$  correspond to  $DN, SN$ , the disc bundle and sphere bundle of  $N$  respectively. Define  $\iota(x) \in U_G^{-n}$  to be the element represented

$$V^+ \xrightarrow{(1)} V^+ / (V^+ \setminus \text{int } T) \cong T / \partial T \cong DN / SN \xrightarrow{(2)} M \vee \mathbb{R}^{-n/2} (G)$$

where (1) is the collapsing map (Thom construction) and (2) is induced by the  $G$ -unitary structure on  $M$ . If  $n$  is odd, embed in  $V \oplus \mathbb{R}$  ( $\mathbb{R}$  the reals with trivial  $G$ -action) and carry out the same construction.

Lemma 4.1.1:

$$\iota : \mathcal{U}_G^* \longrightarrow U_G^*$$

is well-defined.

Proof:

(i)  $\iota$  does not depend on the choice of  $V$ :

Suppose  $f : M \rightarrow V, f' : M \rightarrow V'$  are two permissible embeddings of  $M$ . There is the diagonal embedding:  
 $d : M \rightarrow V \oplus V'$  given by  $d(m) = f(m) \oplus f'(m)$ .

We need only show that  $f$  and  $d$  give the same answer for

$\dot{i}([M])$ . Consider the  $G$ -homotopy of embeddings  $M \rightarrow V \oplus V'$

defined by  $d_t(m) = f(m) + t f'(m)$   $[0 \leq t \leq 1]$ . This is

a  $G$ -homotopy between  $d_0$  and  $d_1 = d$ . Since  $\dot{i}$  depends only on the  $G$ -homotopy class, it is enough to compare  $f: M \rightarrow V$

and  $d_0: M \rightarrow V \oplus V'$ . Let  $N$  be the normal bundle of  $f(M)$ .

Then  $N \oplus V'$  is the normal bundle of  $d_0(M)$  in  $V \oplus V'$ . Hence

it is easily seen that the representative of  $\dot{i}(x)$  given by

considering the embedding  $d_0: M \rightarrow V \oplus V'$  is given by

suspending the  $G$ -map representing  $\dot{i}(x)$ , (using  $f: M \rightarrow V$ ),

by  $V'^+$ . This completes the proof of (i).

(ii)  $\dot{i}$  does not depend on the representative of the bordism

class : see C - F [7] P.28 and tom Dieck [11] P33, Q.E.D.

It has been shown (Thom-Milnor [23] and [17]) that in the

ordinary case  $\dot{i}: U^* = \mathcal{H}_e^* \rightarrow U^*$  is an isomorphism.

The inverse homomorphism is defined by using:

#### Th 4.1.2:

Given a base point preserving map:

$$f: S^{n+k} \longrightarrow M_k,$$

$\exists$  a based map  $h: S^{n+k} \longrightarrow M_k$  s.t.

(i)  $h$  is homotopic to  $f$ , the homotopy fixed at the base points,

(ii)  $h$  is differentiable and (iii)  $h$  is transverse regular to

$$B_k = B_k(e)$$

(Proved as in [15]).



Remark 4.1.3:

For  $G$  abelian,  $G \neq e$  the map:

$$i: \mathcal{U}_G^* \longrightarrow U_G^*$$

is not an isomorphism.

Proof:

Let  $V$  be a non-trivial irreducible representation of  $G$ . Thus it is one-dimensional. The space  $M_1(G)$  can be naturally identified with  $P(V^{\infty} \oplus 1)$  = the projective space of  $V^{\infty} \oplus 1$  (see Lemma 2.1.3) and so we can define a based  $G$ -map:

$$P(V)^+ \xrightarrow{f} M_1(G)$$

by sending  $P(V) = \text{point} \longrightarrow V \subset M_1(G)$ . It is unique up to  $G$ -homotopy (Proposition 1.3.1). Let  $\rho \in U_G^2(\text{point})$  be the element represented by this map. By Cor 2.1.6,  $\bar{\lambda}^{-1}$  is represented by  $1 - H \in \tilde{K}_G(P(W \oplus 1)) \forall G\text{-module } W, H$  being the Hopf bundle over  $P(W \oplus 1)$ . It follows that  $\mu(\rho) = 1 - V \neq 0$  in  $K_G(\text{pt}) = RG$ , since the latter is the free group generated by the irreducible, inequivalent representations of  $G$ . Hence  $U_G^2$  has non-zero elements e.g.  $\rho$ . But since  $\mathcal{U}_G^n = 0$  for all  $n > 0$ , then  $i: \mathcal{U}_G^* \longrightarrow U_G^*$  is not an isomorphism.

Q.E.D.

Remark 4.1.4:

The equivariant analogue of Th 4.1.2 is false (at least for  $G$  abelian,  $G \neq e$ ).

Proof:

Suppose otherwise. Let  $P(V)^+ \xrightarrow{f} M_1(G)$  be the  $G$ -map we have defined in the proof of Remark 4.1.3. Then  $\exists$  a  $G$ -module  $W$  such that  $P(V)^+ \xrightarrow{f} M_1(G)$  can be factored in the form

$$P(V)^+ \xrightarrow{j} P(W \oplus 1) \subset M_1(G)$$
 and  $j$  is transverse regular on  $P(W) \subset P(W \oplus 1)$ . This implies that  $j$  assigns to  $P(V)$  a point of  $P(W \oplus 1) - P(W)$ . The latter  $G$ -deformation retracts onto  $P(1)$ . Hence  $f$  is  $G$ -homotopic to the map sending  $P(V)^+$  to the base point of  $M_1(G)$  (by a based homotopy). Hence the element  $\rho \in U_G^2$  represented by  $f$  is zero - contradiction (see the previous proof). Q.E.D.

#### § 4.2 : A Gysin homomorphism:

In this section we assume that  $G$  is a compact connected Lie group. Let  $T$  be a maximal torus of  $G$ .  $G/T$  is a complex  $G$ -manifold which admits a complex  $G$  embedding in a  $G$ -module  $V$ . In the usual way, the normal bundle of  $G/T$  in  $V$  can be identified with a tubular neighbourhood  $N$  of it. Let  $V^+ \xrightarrow{f} N^+$  be the collapsing map (Thom construction) and let  $N^+ \xrightarrow{h} M_k(G)$  be induced by a classifying map of the complex  $G$ -vector bundle  $N \rightarrow G/T$ . As pointed out in § 4.1, the composition :  $V^+ \xrightarrow{f} N^+ \xrightarrow{h} M_k(G)$  is a representative of  $i([G/T])$  where  $i : \mathcal{U}_G^* \rightarrow \mathcal{U}_G^*$  is the natural map. Suppose now  $X$  is a compact  $G$ -space. Let

$$\phi : U_G^* (G/T \times X) \longrightarrow \tilde{U}_G^* (N^+ \wedge X^+), \quad \phi' : U_G^* (X) \longrightarrow \tilde{U}_G^* (V^+ \wedge X^+)$$

be the Thom isomorphisms (Th 1.4.5). Define

$$P_* : U_G^* (G/T \times X) \longrightarrow U_G^* (X)$$

to be equal to the composition:

$$U_G^* (G/T \times X) \xrightarrow{\phi} \tilde{U}_G^* (N^+ \wedge X^+) \xrightarrow{f^* \wedge 1} \tilde{U}_G^* (V^+ \wedge X^+) \xrightarrow{\phi'^{-1}} U_G^* (X).$$

Lemma 4.2.1:

$P_*$  is a well-defined homomorphism.

Proof:

One needs to verify that  $P_*$  does not depend on the choice of the embedding of  $G/T$ . We use an Atiyah type of argument

(cf [5] 498-499). As in the proof of Lemma 4.1.1, it is

enough to compare the results of embeddings  $S: G/T \longrightarrow V$  and

$d_0: G/T \longrightarrow V \oplus V'$  such that if  $N$  is the normal bundle of  $S$  in  $V$ , then  $N \oplus V'$  is the normal bundle of  $d_0$  in  $V \oplus V'$ .

Denote by  $S_*$  the composition

$$U_G^* (G/T \times X) \xrightarrow{\phi} \tilde{U}_G^* (N^+ \wedge X^+) \xrightarrow{f^* \wedge 1} \tilde{U}_G^* (V^+ \wedge X^+)$$

and by  $d_{0*}$  the composition

$$U_G^* (G/T \times X) \xrightarrow{\phi} \tilde{U}_G^* ((N \oplus V')^+ \wedge X^+) \xrightarrow{f^* \wedge 1} \tilde{U}_G^* ((V \oplus V')^+ \wedge X^+).$$

Because of the transitivity of the Thom homomorphism, the diagram

$$\begin{array}{ccc} U_G^* (G/T \times X) & & \\ \downarrow S_* & \searrow d_{0*} & \\ \tilde{U}_G^* (V^+ \wedge X^+) & \xrightarrow{\phi} & \tilde{U}_G^* ((V \oplus V')^+ \wedge X^+) \\ \uparrow \phi' & \nwarrow \phi & \\ & U_G^* (X) & \end{array}$$

is commutative. Hence the lemma.

Q.E.D.

Let  $p^* : U_G^*(X) \longrightarrow U_G^*(G/T \times X)$  be the natural homomorphism induced by the projection  $G/T \times X \longrightarrow X$ .

Proposition 4.2.2:

Given a compact  $G$ -space  $X$ , then for all  $x \in U_G^*(G/T \times X)$ , and  $y \in U_G^*(X)$ ,  $p_*(x \cdot p^*(y)) = p_*(x) \cdot y$ . Moreover the composition  $U_G^*(X) \xrightarrow{p^*} U_G^*(G/T \times X) \xrightarrow{p_*} U_G^*(X)$  is multiplication by the  $G$ -bordism class  $[G/T] \in \mathcal{H}_G^*$ .

Proof:

The first part follows by the method of Dyer [12] P54. Since  $X$  is compact,  $U_G^*(G/T \times X)$  has an identity 1, so that  $p_* \circ p^*(y) = p_*(1) \cdot y$ . The naturality of  $p_*$  gives rise to a commutative diagram:

$$\begin{array}{ccc} U_G^*(G/T) & \xrightarrow{p_*} & U_G^* \\ \downarrow k^* & & \downarrow k^* \\ U_G^*(G/T \times X) & \xrightarrow{p_*} & U_G^*(X) \end{array}$$

where  $k^*$  is induced by  $X \longrightarrow \text{point}$ . Hence  $p_*(1) = k^*(p_*(1))$ .

Going back to the first paragraph of this section, we see that  $p^*(1)$  is represented by the composition  $V^+ \xrightarrow{f} N^+ \xrightarrow{h} M_k(G)$  i.e.  $p^*(1) = ([G/T])$ . Q.E.D.

By Lemma 1.4.7.  $p^* : U_G^*(X) \longrightarrow U_G^*(G/T \times X)$  can be factored as the composition:

$$U_G^*(X) \xrightarrow{\Gamma} U_T^*(X) \xrightarrow{F(T,G)} U_G^*(G \times_T X) \xrightarrow{v^{*-1}} U_G^*(G/T \times X).$$

Define  $\tau_! : U_T^*(X) \longrightarrow U_G^*(X)$  by the composition:

$$U_T^*(X) \xrightarrow{F(T,G)} U_G^*(G \times_T X) \xrightarrow{v^* - 1} U_G^*(G/T \times X) \xrightarrow{P_*} U_G^*(X).$$

We summarize the results of this section in

Cor 4.2.3:

Let  $r : U_G^*(X) \longrightarrow U_T^*(X)$  be the restriction homomorphism ( $X$  compact). There is a natural homomorphism

$$\tau_! : U_T^*(X) \longrightarrow U_G^*(X)$$

such that  $\tau_! \circ r$  is multiplication by  $[G/T]$ .

4.3 : The Equivariant Todd Genus of  $G/T$  :

We assume the following from [4] sections 6.2, 6.3 (see also [5]).

Th 4.3.1:

Let  $\tau^{G/T}$  be the tangent bundle of  $G/T$ ,  $\varphi : K_G(G/T) \longrightarrow K_G(\tau^{G/T})$  the Thom homomorphism, and  $t\text{-ind} : K_G(\tau^{G/T}) \longrightarrow RG$  the topological index. Then  $t\text{-ind}(\varphi(1)) = 1$ .

This implies

Proposition 4.3.2:

The composition  $\mathcal{U}_G^* \xrightarrow{i} U_G^* \xrightarrow{\mu} RG$  assigns to  $[G/T]$  the identity of  $RG$ .

Proof:

There is a complex embedding of  $G/T$  in a  $G$ -module  $V$ . In the usual way, the normal bundle  $N$  to  $G/T$  in  $V$  can be identified with a  $G$ -tubular neighbourhood of it. There is then induced a complex embedding of  $G/T$  in  $\tau V$  with normal bundle  $\tau N$ . Let  $(\tau V)^+ \xrightarrow{k} (\tau N)^+$  be the collapsing map and let  $(\tau N)^+ \xrightarrow{+f} M_r(G)$

be induced by a classifying map of  $\tau_N \rightarrow G/T$  ( $r = 21V1 - 1 \cdot G/T \cdot 1$ ).  
 By the definition of  $i: \mathcal{H}_G^* \rightarrow U_G^*$ ,  $i([G/T])$  is represented  
 by the composition:

$(\tau V)^+ \xrightarrow{k} (\tau N)^+ \xrightarrow{f} M_r(G)$ . Hence  $\mu i([G/T])$  is the  
 image of the natural element  $\bar{\lambda}^r$  in  $\tilde{O}(M_r(G))$  under the  
 composition:

$$\tilde{O}(M_r(G)) \xrightarrow{f!} \tilde{K}_G((\tau N)^+) \xrightarrow{k!} \tilde{K}_G((\tau V)^+) \xrightarrow{\Phi^{-1}} RG.$$

By naturality of the Thom class,  $f!(\bar{\lambda}^r) =$  the Thom class in  
 $\tilde{K}_G((\tau N)^+)$  of the bundle  $\tau_N \rightarrow G/T$ . Denoting the Thom  
 homomorphism by  $\Phi$ , we deduce that applying the composition:

$$K_G(G/T) \xrightarrow{\Phi} K_G(\tau N) \xrightarrow{k!} K_G(\tau V) \xrightarrow{\Phi^{-1}} RG$$

to the element  $1 \in K_G(G/T)$  gives us  $\mu i([G/T])$ . Let  $E_x$   
 denote the fiber over the point  $x$  of the bundle  $E \rightarrow X$ .

Given  $x \in G/T$ ,  $(\tau N)_x = N_x \oplus V = N_x \oplus N \oplus (\tau^{G/T})_x$ .

Hence  $\tau^{G/T} \rightarrow G/T$  is naturally a  $G$ -sub bundle of  $\tau_N \rightarrow G/T$   
 (when both are considered as complex bundles). By the

transitivity of the Thom homomorphism ([20]), we get a  
 commutative diagram:

$$\begin{array}{ccccc} K_G(G/T) & \xrightarrow{\Phi} & K_G(\tau N) & \xrightarrow{k!} & K_G(\tau V) \xrightarrow{\Phi^{-1}} RG \\ & \searrow \Phi & \uparrow \Phi & \nearrow \Phi & \\ & & K_G(\tau^{G/T}) & & \end{array}$$

Therefore,  $\mu i([G/T]) = t\text{-ind}(\Phi(1)) = 1$  (Theorem 4.3.1).

Q.E.D.

Remark:

Conner-Floyd have proved that for ordinary unitary bordism, the composition  $\mathcal{U}_* \xrightarrow{i} U^* \xrightarrow{\mu} \mathbb{Z}$  sends the bordism class of a manifold  $M^{2n}$  into  $(-)^n \text{Td}[M^{2n}]$  where  $\text{Td}[M^{2n}]$  is the Todd genus of  $M^{2n}$  ([8] P.37). So we view the above proposition as a computation (up to sign) of the equivariant Todd genus of  $G/T$ .

§ 4.4: The Main Theorem:

Proceeding as in § 4.2, we can construct a homomorphism

$$P! : K_G^* (G/T \times X) \longrightarrow K_G^* (X)$$

for all compact  $G$ -spaces  $X$  ( $G$  connected). Let  $r' : K_G^* (X) \rightarrow K_T^* (X)$  be the restriction homomorphism, and let  $K_T^* (X) \xrightarrow{F'(T,G)} K_G^* (G \times_T X)$

be the canonical isomorphism ([20] P.132). Using the identification  $v : G \times_T X = G/T \times X$  (1.4.6), we get an isomorphism

$$v! : K_G^* (G/T \times X) \stackrel{\sim}{=} K_G^* (G \times_T X).$$

Lemma 4.4.1:

The diagrams:

$$(i) \quad \begin{array}{ccc} U_G^* (X) & \xrightarrow{r} & U_T^* (X) \\ \mu \downarrow & & \downarrow \mu \\ K_G^* (X) & \xrightarrow{r'} & K_T^* (X) \end{array}$$

$$(ii) \quad \begin{array}{ccc} U_T^* (X) & \xrightarrow{F(T,G)} & U_G^* (G \times_T X) \\ \mu \downarrow & & \downarrow \mu \\ K_T^* (X) & \xrightarrow{F'(T,G)} & K_G^* (G \times_T X) \end{array}$$

$$(iii) \quad \begin{array}{ccc} U_G^* (G/T \times X) & \xrightarrow{P_*} & U_G^* (X) \\ \mu \downarrow & & \downarrow \mu \\ K_G^* (G/T \times X) & \xrightarrow{P!} & K_G^* (X) \end{array}$$

are commutative.

Proof:

The proof of (i) is straightforward. To prove (ii) is commutative, we recall that the inverse isomorphisms

$$F(G,T) : U_G^*(G \times X) \longrightarrow U_T^*(X) \quad \text{and} \quad F'(G,T) : K_G^*(G \times X) \longrightarrow K_T^*(X)$$

are given by the compositions  $U_G^*(G \times X) \xrightarrow{r} U_T^*(G \times X) \xrightarrow{q^*} U_T^*(X)$

and  $K_G^*(G \times X) \xrightarrow{r} K_T^*(G \times X) \xrightarrow{q^*} K_T^*(X)$  where  $r$  is the restriction homomorphism and  $q : X \rightarrow G \times X$  sends  $x$  to  $[1, x]$ .

Hence by (i) and the naturality of  $\mu$ , (ii) is commutative.

Because  $\mu$  is natural and commutes with the Thom homomorphism (Lemma 2.1.7), (iii) is commutative. Q.E.D.

Define  $r'_! : K_T^*(X) \longrightarrow K_G^*(X)$  by the composition

$$K_T^*(X) \xrightarrow{F(T,G)} K_G^*(G \times X) \xrightarrow{v!^{-1}} K_G^*(G/T \times X) \xrightarrow{P!} K_G^*(X).$$

Hence  $r'_! \circ r^* = \text{id} : K_G^*(X) \longrightarrow K_G^*(X)$  (Atiyah [2]). On the other hand, the composition

$$(4.4.2) \quad U_G^*(X) \otimes_{U_G}^* RG \xrightarrow{r \otimes r^*} U_T^*(X) \otimes_{U_T}^* RT \xrightarrow{r_! \otimes r'_!} U_G^*(X) \otimes_{U_G}^* RG$$

is multiplication by  $\mu([G/T]) = 1$  (Cor 4.2.3 and Prop. 4.3.2) i.e.

the composition (4.4.2) is equal to  $\text{id}$ :

$$U_G^*(X) \otimes_{U_G}^* RG \longrightarrow U_G^*(X) \otimes_{U_G}^* RG.$$

Combining this result with Th. 3.4.2 and Lemma 4.4.1.

Theorem 4.4.3:

Let  $G$  be a compact connected Lie group. The homomorphism

$$\mu' \otimes 1 : U_G^*(X) \otimes_{U_G}^* RG \longrightarrow K_G^*(X)$$



is an isomorphism of  $\mathbb{Z}_2$ -graded rings for all compact  $G$ -spaces  $X$ . Moreover for any compact Lie group  $G$ , and a compact  $G$ -space  $X$ ,

$$\mu \otimes 1 : U_G^*(X) \otimes_{U_G^*} RG \cong K_G^*(X).$$

Proof:

Case (i) :  $G$  connected. Let  $T$  be a maximal torus of  $G$ .

By Theorem 3.4.2,  $\mu \otimes 1 : U_T^*(X) \otimes_{U_T^*} RT \cong K_T^*(X)$ . From Lemma 4.4.1, the diagram:

$$\begin{array}{ccccc} U_G^*(X) \otimes_{U_G^*} RG & \xrightarrow{r \otimes r'} & U_T^*(X) \otimes_{U_T^*} RT & \xrightarrow{r_! \otimes r'_!} & U_G^*(X) \otimes_{U_G^*} RG \\ \downarrow \mu \otimes 1 & & \downarrow \mu \otimes 1 & & \downarrow \mu \otimes 1 \\ K_G^*(X) & \xrightarrow{r'} & K_T^*(X) & \xrightarrow{r'_!} & K_G^*(X) \end{array}$$

is commutative. Since  $\mu \otimes 1$  and  $r \otimes r'$  are injective (4.4.2), then  $\mu \otimes 1$  is injective. It is also surjective because  $r_!$  and  $\mu \otimes 1$  are. Hence it is an isomorphism.

Case (ii) : The general case. Embed  $G$  in  $U(n)$  for some  $n$ .

$$\begin{aligned} \text{Then } U_G^*(X) \otimes_{U_G^*} RG &\cong U_{U(n)}^*(U_{n/G} \times X) \otimes_{U_G^*} K_{U(n)}^*(U_n/G) \cong \\ &\cong U_{U(n)}^*(U_{n/G} \times X) \otimes_{U_G^*} (U_{U(n)}^*(U_n/G) \otimes_{U_n^*} RU_n) \\ &\cong (U_{U(n)}^*(U_{n/G} \times X) \otimes_{U_G^*} U_G^*) \otimes_{U_n^*} RU_n \cong U_{U(n)}^*(U_{n/G} \times X) \otimes_{U_n^*} RU_n \\ &\cong K_{U(n)}^*(U_{n/G} \times X) \cong K_G^*(X) \text{ where all the isomorphisms that} \\ &\text{appear are the canonical ones we already defined (see §1.4, and} \\ &\text{case (i) above).} \end{aligned}$$

REFERENCES

1. M.F. Atiyah , K-theory (Benjamin 1967)
2. On Both Periodicity, The Quarterly Journal of Maths  
Vol. 19, No. 74, June 1968
3. Bordism and Cobordism, Cambridge Phil.Soc. Vol.57, 1961
4. M.F. Atiyah, and G.B. Segal,  $K_G$  - theory Notes (Oxford) 1965
5. M.F. Atiyah and I.M. Singer, The index of Elliptic operators, Ann.  
of Math 87 (1968)
6. N. Bourbaki, 'Algebre commutative'. Ch. 2. Localization, Herman,  
Paris (1961)
7. P.E. Conner and E.E. Floyd, 'Differentiable Periodic Maps'  
Springer (1964)
8. , 'The relation of cobordism to K-theory'  
Lecture notes in Math. 28, Springer 1966
9. , 'Torsion in SU-bordism' Mem. Amer.  
Math. Soc. 60 (1966)
10. H. Cartan and S. Eilenberg, Homological Algebra, Princeton Univ.  
Press (1956)
11. T. tom Dieck, 'Cobordism Theory and Transformation G maps'  
Aarhus Universitet (1968/69) No. 30
12. E. Dyer 'Cohomology Theories'. Benjamin (1969)
13. S.G.Hoggar 'On  $KO$ -theory of Grassmannians Ph.D thesis  
Warwick University
14. D. Husemoller 'Fibre Bundles' McGraw-Hill
15. J.W. Milnor 'Characteristic Classes' (1957)
16. 'Differential Topology Notes' Lect.Notes (1958)
17. 'On the cobordism ring....' Amer. J. Math. 82 (1960)
18. R.S. Palais, 'Imbedding of compact, diff. transformation groups',  
J. Math. Mech 6 (1957)
19. Pontrjagin, Lie Groups
20. G.B. Segal, Equivariant K-theory, Publ. Math I. H.E.S. (1968)
21. , Classifying spaces, Publ. math I. H.E.S. (1968)
22. Spanier, Algebraic Topology
23. R. Thom, Quelques proprietes globales des varietes diff. Comment Math.  
Helv. 27 (1953)
24. G. Whitehead, 'Generalized homology theories' Trans. Amer. Math. Soc.  
102, (1962)